



Marine oxygenation, lithistid sponges, and the early history of Paleozoic skeletal reefs

Jeong-Hyun Lee^{a,*}, Robert Riding^b

^a Department of Geology and Earth Environmental Sciences, Chungnam National University, Daejeon 34134, Republic of Korea

^b Department of Earth and Planetary Sciences, University of Tennessee, Knoxville, TN 37996, USA

ARTICLE INFO

Keywords:

Cambrian
Reef gap
Dysoxia
Hypoxia
Lithistid sponge-microbial reef

ABSTRACT

Microbial carbonates were major components of early Paleozoic reefs until coral-stromatoporoid-bryozoan reefs appeared in the mid-Ordovician. Microbial reefs were augmented by archaeocyath sponges for ~15 Myr in the early Cambrian, by lithistid sponges for the remaining ~25 Myr of the Cambrian, and then by lithistid, calathiid and pulchrlaminid sponges for the first ~25 Myr of the Ordovician. The factors responsible for mid-late Cambrian microbial-lithistid sponge reef dominance remain unclear. Although oxygen increase appears to have significantly contributed to the early Cambrian 'Explosion' of marine animal life, it was followed by a prolonged period dominated by 'greenhouse' conditions, as sea-level rose and CO₂ increased. The mid-late Cambrian was unusually warm, and these elevated temperatures can be expected to have lowered oxygen solubility, and to have promoted widespread thermal stratification resulting in marine dysoxia and hypoxia. Greenhouse conditions would also have stimulated carbonate platform development, locally further limiting shallow-water circulation. Low marine oxygenation has been linked to episodic extinctions of phytoplankton, trilobites and other metazoans during the mid-late Cambrian. We propose that this tendency to dysoxia-hypoxia in shallow marine environments also limited many metazoan reef-builders. In contrast, during the mid-late Cambrian, the ability of lithistid sponges to withstand low oxygen levels allowed them to create a benthic association with microbial carbonates that dominated global reefs. These conditions ameliorated during the Ordovician, as temperature decline promoted ocean ventilation. The prolonged time gap occupied by low diversity reefs between the 'Cambrian Explosion' and the 'Great Ordovician Biodiversification Event' reflects elevated temperatures and reduced marine oxygenation that limited metazoan diversification in shallow marine environments.

1. Introduction

The early Paleozoic radiation of marine animals, commonly referred to as the 'Cambrian Explosion' (Cloud, 1948; Bowring et al., 1993; Conway Morris, 2006), stalled in the mid-Cambrian and did not regain impetus until the mid-Ordovician (Sepkoski Jr., 1981; Webby, 2002; Bambach et al., 2004) during the 'Great Ordovician Biodiversification Event' (Webby et al., 2004). Reef development shows a similar pattern, leading to the observation that mid-late Cambrian reef communities were dominantly microbial (Fagerstrom, 1987; Zhuravlev, 1996; Rowland and Shapiro, 2002). As a result, the prolonged interval between late early Cambrian extinction of archaeocyath sponges and the Early Ordovician rise of lithistid sponge-microbial-calathiid reef communities came to be regarded as a "reef gap" (Sheehan, 1985; Zhuravlev, 1996; Rowland and Shapiro, 2002; Kiessling, 2009). Challenges to this view first emerged in the report of early Furongian (late Cambrian) lithistid sponge-microbial (LSM) reefs in Iran (Hamdi et al.,

1995). Additional examples of mid-late Cambrian LSM reefs have since been recognized throughout Laurentia and Gondwana (Mrozek et al., 2003; Shapiro and Rigby, 2004; Johns et al., 2007; Kruse and Zhuravlev, 2008; Hong et al., 2012, 2016; Kruse and Reitner, 2014; Adachi et al., 2015; Lee et al., 2016a). It is now clear that LSM reef communities were globally distributed for ~25 Myr during the mid-late Cambrian. This community was augmented in the Early Ordovician by a variety of skeletal organisms, such as *Calathium*, *Pulchrlamina*, *Amassia*, bryozoans, stromatoporoids, and *Lichenaria*, but reef diversification remained relatively slow until bryozoans, corals and stromatoporoids became common in the mid-Ordovician, ~460 Ma (Fagerstrom, 1987; Webby, 2002; Servais et al., 2009). Numerical ages used here accord with those of the International Chronostratigraphic Chart (Cohen et al., 2013). Recognition of mid-late Cambrian LSM reefs closes the "reef gap" (Lee et al., 2016a and references therein) but also raises questions. What factors contributed to lithistid success following archaeocyath decline, why did LSM reefs dominate the remainder of the

* Corresponding author.

E-mail addresses: jeonghyunlee@cnu.ac.kr (J.-H. Lee), rriding@utk.edu (R. Riding).

Cambrian, and why did it take so long for the bryozoan-coral-stromatoporoid community to emerge in the Ordovician? Explanations for these aspects of reef developments could shed light on broader patterns of change that continue to be debated, particularly the seemingly slow overall pace of mid–late Cambrian animal diversification (Sepkoski Jr., 1981; Pruss et al., 2010; Knoll and Fischer, 2011; McKenzie et al., 2014; Saltzman et al., 2015).

Here we explore the possibility that prolonged low levels of marine oxygenation significantly influenced the relative success and development of reef communities during the mid–late Cambrian and Early Ordovician (Webby, 2002; Saltzman et al., 2015; Lee and Riding, 2016). High mid–late Cambrian sea-level and elevated atmospheric CO₂ promoted ‘greenhouse’ conditions with elevated temperatures (Berner, 1990, 2009; McKenzie et al., 2014). Higher temperatures decrease oxygen solubility in seawater, increase thermal stratification of the ocean, and favor marine productivity, while simultaneously stimulating carbonate platform development that restricts shallow-water circulation. Collectively, these factors tend to induce episodic marine dysoxia-hypoxia (Saltzman, 2005; Gill et al., 2011) that has been linked to repeated extinctions of phytoplankton, trilobites and other metazoans during the mid–late Cambrian (e.g., Bambach et al., 2004).

We propose that lithistid sponges, along with microbes, were predisposed to tolerate low oxygen levels, and that this was a central factor in their development and persistence during the mid–late Cambrian. These conditions began to change in the Early Ordovician as temperature decline (Trotter et al., 2008) promoted ocean ventilation (Thompson et al., 2012; Kah et al., 2016; Young et al., 2016). In the mid-Ordovician the bryozoan-coral-stromatoporoid community established itself as the next major phase of Paleozoic reef development (Pitcher, 1964; Fagerstrom, 1987). From this perspective, the relationship between reef community evolution and dysoxia-hypoxia is a key to understanding not only the Cambrian “reef gap”, but also the relatively slow progress of animal radiation in mid-Cambrian to mid-Ordovician shallow marine environments.

2. Cambrian–Ordovician marine diversity

Marine animal generic diversity shows a plateau in the late Cambrian (Furongian Epoch; Paibian and Jiangshanian ages) (~510–485 Ma) (Sepkoski Jr., 1979, 1981), before recovering near the Cambrian–Ordovician transition (Na and Kiessling, 2015). This interval of depressed diversity largely reflects archaeocyath extinctions, followed by trilobite extinctions and low levels of origination (Bambach et al., 2004). Mid–late Cambrian and Early Ordovician trilobite extinctions define “biomere” boundaries (Palmer, 1984; Taylor, 2006) that, as in the case of the late Cambrian Age 3 Sinsk Event (Zhuravlev and Wood, 1996), have been linked to influxes of anoxic and/or cold basinal waters onto shallow shelves (Stitt, 1975; Palmer, 1984; Westrop and Ludvigsen, 1987). This is consistent with survival of trilobites adapted to reduced oxygen availability at biomere boundaries (Palmer, 1984; Taylor, 2006; McKenzie et al., 2014; Saltzman et al., 2015). In addition, dysoxia-tolerant arthropod groups such as olenids (Fortey, 2000) and phosphatocopids (Williams et al., 2011) increased in diversity in the late Cambrian (Fig. 1).

Biomere and other Cambrian biotic turnovers/extinctions have in turn been linked to carbon isotope excursions (Taylor, 2006; Zhu et al., 2006; Peng et al., 2012; Saltzman et al., 2015). In addition to the ~510 Ma Sinsk Event, these include the early Paibian Marju-miid–Pteroccephaliid biomere boundary and the early Jiangshanian Pteroccephaliid–Ptychaspis biomere boundary, which approximately correspond with the lower and upper boundaries respectively of the Steptoean positive carbon isotope excursion (SPICE) event (Saltzman et al., 1995, 2000; Perfetta et al., 1999; Saltzman, 1999; Montañez et al., 2000; Lee et al., 2015a; Gerhardt and Gill, 2016) (for correlations see Taylor et al., 2012, fig. 1) (Fig. 1). We infer, along with others, that mid–late Cambrian extinctions occurred as poorly oxygenated basinal

water encroached onto carbonate platforms, stressing shallow-water communities (Palmer, 1984; Berry et al., 1989; Taylor, 2006; Gill et al., 2011; Landing, 2012a, 2012b; Saltzman et al., 2015). Similar effects have been invoked to account for extinction of carbonate platform communities in Jurassic–Cretaceous Oceanic Anoxic Events (Schlanger and Jenkyns, 1976; Jenkyns, 1991, 2010; Parente et al., 2008).

Marine diversity greatly increased during the Ordovician (Sepkoski Jr., 1981). The Great Ordovician Biodiversification Event (GOBE) (Webby et al., 2004) was a diachronous and extended event, initiated in the Furongian and accelerating in the Darriwilian, that continued until almost the end of the Ordovician (Sepkoski Jr., 1981; Webby et al., 2004; Harper, 2006; Servais et al., 2010, 2016). Significant increases in animal biodiversity at family, genus and species level resulted in the rise of the ‘Paleozoic Evolutionary Fauna’, dominated by suspension-feeding organisms such as brachiopods, rugose and tabulate corals, crinoids, bryozoans, and stromatoporoids (Sepkoski Jr., 1979; Bambach et al., 2002). This community developed throughout the Early Ordovician, and by the late Darriwilian had a radical affect on reef building (Pitcher, 1964; Fagerstrom, 1987; Webby, 2002; Webby et al., 2004). In addition, global increase in phytoplankton (acritarch) diversity, broadly paralleling that of animals, occurred in the Furongian and Early Ordovician (Servais et al., 2008, 2016; Nowak et al., 2015). These Ordovician upturns in diversity have been broadly related to increase in atmospheric O₂ (Servais et al., 2008, 2010, 2016; Saltzman et al., 2011; Edwards et al., 2017) and decline in global temperature (Trotter et al., 2008).

3. Late Ediacaran–mid-Ordovician eukaryote reef development

Microbial carbonates (stromatolites, thrombolites and associated calcimicrobes) generally dominated reefs from the Ediacaran to the Early Ordovician. The earliest skeletal reefs involved Ediacaran calcified sessile benthic eukaryotes (*Cloudina*, *Namacalathus*, *Namapoikia*) and early Cambrian “*Ladatheca*”. Archaeocyaths, and locally radiocyaths and coralomorphs, are conspicuous in early Cambrian reefs. Archaeocyaths and coralomorphs continued – although much reduced – during the mid–late Cambrian, when lithistid sponges were the most prominent components of skeletal reefs. Large lithistids (*Archaeoscyphia*) appeared in the Early Ordovician, augmented by calathiids, pulchrlaminids, and the first undisputed tabulate corals, bryozoans and stromatoporoids. These latter appearances set the scene for the mid-Ordovician diversification of stromatoporoids, tabulates and bryozoans, that created reefs dominated by encrusting and massive skeletons at the end of the Darriwilian.

The affinities of many Ediacaran–Ordovician skeletons remain problematic. *Namapoikia* may be a chaetetic sponge or simple colonial cnidarian (Wood et al., 2002; Wood, 2017). Archaeocyaths and lithistids are sponges. Radiocyath skeletal structure resembles that of *Calathium*, which is sponge-like (Church, 1991; Rowland, 2001), and also that of cyclocrinids (Beadle, 1988); but these groups have also been compared with green algae (e.g., Nitecki and Debrenne, 1979). Coralomorphs comprise a heterogeneous semi-formal group encompassing massive as well as cuplike skeletons; some of them are considered real corals (e.g., Scrutton, 1997; Hicks, 2006). Pulchrlaminids are stromatoporoid-like, but regarded as a separate group of hypercalcified sponges (Webby, 2015).

Based on their eukaryote skeletal components, we distinguish six reef intervals during this ~100 Myr period (~550–444 Ma), when reefs evolved from mainly microbial into mainly skeletal structures. Time scales are based on Peng et al. (2012), Cohen et al. (2013), and Ogg et al. (2016). These main reef intervals are: I, late Ediacaran (~550–541 Ma): microbial-*Cloudina*; II, Fortunian–mid-Age 2 (541–525 Ma): microbial-“*Ladatheca*”; III, mid-Age 2/late Age 4 (525–510 Ma): microbial-archaeocyath; IV, late Age 4/end Age 10 (510–485 Ma): microbial-lithistid; V, Tremadocian–middle Darriwilian (485–460 Ma): microbial-lithistid-calathiid-pulchrlaminid; VI, late Darriwilian–end-Ordovician (460 Ma–444 Ma): stromatoporoid-bryozoan-tabulate-receptaculitid-microbial.




Fig. 1. Cambrian to Middle Ordovician oxygen level indicators and associated changes in biotic diversity. Generic diversity of calcifying animal groups (Anthozoa, Archaeocyathida, Articulata, Bryozoa, Demospongia, Echinodermata, Mollusca, Ostracoda, and Trilobita) after Pruss et al. (2010). Olenid trilobite diversity from <http://fossilworks.org/> (accessed 22nd Dec. 2017). Bivalved arthropod phosphatocoid diversity from Williams et al. (2011). Extinction events adapted from: 1. Zhuravlev and Wood (1996), 2. Palmer and James (1979), 3. Peng et al. (2012), 4. Palmer (1984), 5. Loch et al. (1993), 6. Adrain et al. (2009). Examples of black shale deposits from Babcock et al. (2015): 1. Pestrotsvet Fm., Russia, 2. Spence Shale, Utah and Idaho, USA, 3. Burgess Shale, British Columbia, Canada, 4. Niutitang Fm., South China, 5. Alum Shale, Scandinavia, 6. Huaqiao Fm., South China, 7. White-leaved Oak Shale, England. Global anoxic events: 1. Sink (Zhuravlev and Wood, 1996), 2. Redlichiiid-Olenellid Extinction Carbon isotope Excursion (ROECE) (Zhu et al., 2006; Faggetter et al., 2017), 3. Steptoean Positive Carbon Isotope Excursion (SPICE) (Saltzman et al., 2000), 4. Reducing condition across the Cambro-Ordovician boundary at Green Point GSSP section (Tripathy et al., 2014). Oxygenation trends: 1. Global ocean oxygenation across the Ediacaran–Cambrian boundary indicated by Mo isotopes (Chen et al., 2015), 2. Global ocean deoxygenation indicated by bioturbation, Mo isotopes, TOC and U (Boyle et al., 2014), 3. Atmospheric oxygen levels calculated by Edwards et al. (2017), 4. Ordovician sea-surface temperatures based on conodont thermometry (Trotter et al., 2008), 5. Tremadocian-Floian ocean oxygenation (Saltzman et al., 2015), 6. Early Darriwilian global ocean ventilation (Kah et al., 2016).

3.1. Late Ediacaran (~550–541 Ma): microbial-*Cloudina*

Calcified sessile problematic metazoans contributed to and locally constructed reefs in the late Ediacaran (~550–541 Ma) (Germs, 1972; Conway Morris et al., 1990; Seilacher, 1999; Grotzinger et al., 2000; Hofmann and Mountjoy, 2001; Penny et al., 2014, 2017; Wood, 2017). Masses of narrow conical calcified *Cloudina* tubes (annelid or cnidarian, Vinn and Zatoñ, 2012; Pacheco et al., 2015), individually up to 8 mm wide and 15 cm long, constructed reefs up to 7 m wide and 3 m high (Wood, 2017). *Cloudina* also often occurs in stromatolite and thrombolite reefs, locally in association with stalked cups of the calcified, but

very thin-walled problematic *Namacalathus* (lophophorate? Zhuravlev et al., 2015) up to 35 mm in size (Grotzinger et al., 2000; Hofmann and Mountjoy, 2001; Zhuravlev et al., 2015; Wood, 2017). *Cloudina* and *Namacalathus* locally mutually attached (Penny et al., 2014, 2017). *Namapoikia*, another early metazoan in late Ediacaran microbial reefs, is tabular crustose and chaetetic-like (see 3.3.4. Chaetetics, below) (Wood et al., 2002), composed of millimetric labyrinthine tubules, and in cavities locally forms sheet-like crusts up to 1 m in extent (Wood, 2017). *Cloudina*, *Namacalathus* and *Namapoikia* are not known to have survived into the Cambrian (Amthor et al., 2003).

3.2. Earliest Cambrian, Fortunian/mid-Age 2 (541–525 Ma): microbial-“*Ladatheca*”

Early Cambrian (Terreneuvian) reefs, like those of the Ediacaran, are predominantly stromatolitic and calcimicrobial (Drosdova, 1980; Riding and Voronova, 1984; Kruse et al., 1996). Among the shelly fossils that appeared in the Fortunian (Bengston, 2004), a single example of a ‘worm reef’ formed by operculate tubular “*Ladatheca*”, individually up to 15 cm long and 6 mm wide, occurs in the late Fortunian (Landing, 1993), and is reminiscent of *Cloudina* reefs.

3.3. Early Cambrian, mid-Age 2/late Age 4 (525–510 Ma): microbial-archaeocyath

3.3.1. Archaeocyaths

Archaeocyaths, previously considered a distinct phylum or compared with coelenterates and algae, are now regarded as sponges (Debrenne et al., 2015b). These distinctive and often heavily calcified skeletons, typically conical with porous walls partitioned and supported by septa, appeared in the lower Tommotian Stage of Siberia (mid-Age 2, ~525 Ma, Ogg et al., 2016) in the zone named after the archaeocyath *Nochorycyathus sunnaginicus* (Fig. 2). Most archaeocyath cups are < 15 mm in diameter and several centimeters in height, but examples 1.5 m in size are reported, and some show mutual attachment that favored reef-construction (Hill, 1964; Wood et al., 1993; Debrenne et al., 2015b). They were long regarded as the first animals to attain a global reef-building role (Fagerstrom, 1987; Rowland and Gangloff, 1988; Wood, 1999; Knoll, 2003; Gandin and Debrenne, 2010), and some early

Cambrian reefs are dominated by archaeocyath frameworks (Rowland, 1984; Riding and Zhuravlev, 1995). However, they are usually also closely associated with abundant calcimicrobes (Pratt et al., 2001; Riding, 2001; Lee et al., 2014b) that diversified in the early Cambrian (Riding and Voronova, 1984; Zhuravlev, 1996), and less commonly with coralomorphs and radiocyaths (Jell, 1984; Kruse et al., 1995, 1996; Zhuravlev and Wood, 1995; Zhuravlev, 2001a; Zhuravlev et al., 2015) (Fig. 3). The peak of archaeocyath diversity in Age 3 (~518 Ma) coincided with that of marine animal genera as a whole (Na and Kiessling, 2015), but the group declined rapidly ~510 Ma (Peng et al., 2012). Archaeocyath decline, sometimes considered the first significant extinction of the Phanerozoic (Newell, 1972; Signor, 1992; Zhuravlev, 1996, 2001a; Bambach et al., 2002; Bambach, 2006), was linked to the Hawke's Bay Regression in the lower Toyonian Stage of Siberia (middle Age 4) by Palmer and James (1979), but Zhuravlev and Wood (1996) suggested that the anoxic Sinsk Event, in the Botomian Stage of Siberia (late Age 3), may have been more important in reducing archaeocyath diversity. Hyoliths, inarticulate brachiopods, and trilobites also show reduced levels of originations in the Botomian (Bambach et al., 2004) (Fig. 1).

3.3.2. Cribricyaths, radiocyaths

During Cambrian Epoch 2, the diverse archaeocyath reef community was augmented by other, mostly problematic, calcified skeletal organisms of various shapes and sizes, such as cribricyaths, radiocyaths and coralomorphs. Cribricyatha (Vologdin, 1961) are small (< 1 cm long), locally abundant, calcareous conical Problematica (Zhuravlev, 2001b) that contributed to reef formation (Wood et al., 1993) in the




Fig. 2. Schematic size, shape, and age summary of reef-building skeletal organisms from late Ediacaran to early Late Ordovician. Time scales from Peng et al. (2012), Cohen et al. (2013), and Ogg et al. (2016). Six reef intervals are recognized: I, Late Ediacaran (~550–541 Ma) microbial reefs with *Cloudina*, *Namacalathus*, and *Namapoikia*. II, Earliest Cambrian (Fortunian–mid-Age 2) microbial reefs with rare “*Ladatheca*”. III, Microbial-archaeocyath sponge reefs with radiocyaths, coralomorphs, etc. from mid-Age 2 to Age 4. IV, Microbial-lithistid sponge reefs in the mid-late Cambrian. V, Microbial-lithistid sponge reefs augmented by *Calathium*, pulchrlaminids, bryozoans, *Lichenaria*, *Amsassia*, and *Cystostroma* in the Early and early Middle Ordovician. VI, Skeletal-dominant reefs mainly constructed by stromatoporoids, corals, and bryozoans together with microbial carbonates in the late Darriwilian and Late Ordovician.




Fig. 3. Relative proportions of main reef components during the Cambrian and Early Ordovician. Early Cambrian microbial-archaeocyath reefs from Gandin and Debrenne (2010), Early Ordovician microbial-lithistid-calathiid reefs from Hong et al. (2015), Li et al. (2015) and references therein. Proportions of mid-late Cambrian reef-building components based on examples in Table 1.

early Cambrian (Tommotian–Botomian) (Debrenne et al., 2015a). An early cribricyath is *Leibaella* (late Tommotian, late Age 2) (Wood et al., 1993). Radiocyaths are similar in size and shape to archaeocyaths, with which they were often confused (Kruse et al., 2015), but were distinguished as a distinct group (Radiocyatha) by Debrenne et al. (1970). An early example is *Gonamisporgia* (late Tommotian, late Age 2). Radiocyaths have globular to pear-shaped skeletons, with outer walls composed of distinctive plates (nesasters) and more closely resemble receptaculitids than archaeocyaths (Nitecki and Debrenne, 1979), although receptaculitids are not common until the Ordovician (see Section 3.5.1. Receptaculitids, radiocyaths, cyclocrinids, below). However, the largest skeletal additions to early Cambrian reef communities were locally made by coralomorphs, such as *Flindersipora* (Fuller and Jenkins, 2007).

3.3.3. Early Cambrian coralomorphs

Tabulate-like phosphatic fossils (e.g., *Ramitubus*) occur in the mid-Ediacaran (~585 Ma) Doushantuo Formation (Van Iten et al., 2014), and *Namapoikia* may be a cnidarian (Wood et al., 2002). Coral-like calcified fossils are diverse in the early Cambrian (Zhuravlev et al., 1993; Wrona and Zhuravlev, 1996; Scrutton, 1999; Debrenne and Reitner, 2001; Hicks, 2006; Fuller and Jenkins, 2007), but are difficult to relate to Tabulata or Rugosa, neither of which is confidently recognized until the Ordovician (Scrutton, 1992; Oliver, 1996; Scrutton, 1997). Jell (1984) placed coral-like early Cambrian fossils within the Coralomorpha. This highly heterogeneous semi-formal group encompasses chaetetids and early “tabulates”, with or without septa and tabulae, as well as cuplike skeletons.

Early coralomorphs are best known from calcimicrobial-archaeocyath reefs in the early Cambrian. They exhibit two contrasting morphotypes: small cup-like and large massive domical. Scrutton (1997) recognized two orders of Cambrian zoantharian corals (Tabulaconida, Debrenne et al., 1987; Cothoniida, Oliver and Coates, 1987) and suggested that calcified early–early middle Cambrian corals represent ‘a series of short-lived calcification episodes among contemporary anemones’ (Scrutton, 1999). Middle Tommotian (523 Ma) *Cysticyathus*, previously included in archaeocyaths, is an early cup-like coralomorph in calcimicrobial-archaeocyath reefs (Debrenne and Reitner, 2001, fig. 14.6b). Somewhat younger cup-like forms include Age 5–Drumian

Cambroctoconus (Park et al., 2011; Peel, 2017b), Age 4 *Cothonion* (Peel, 2011) and *Lipopora* (see Park et al., 2016). Some of these formed mid-Cambrian reefs, e.g., *Cambroctoconus* (Lee et al., 2016a), which has been described as a stem-group cnidarian based on its octoral symmetry (Park et al., 2011).

Much larger domical forms in the late Botomian (~513 Ma) include *Flindersipora* (Lafuste et al., 1991; Savarese et al., 1993; Debrenne and Reitner, 2001; Fuller and Jenkins, 2007), *Yaworipora* (Zhuravlev, 1999), *Moorowipora* (Sorauf and Savarese, 1995), *Arrowipora* and *Blinmanipora* (Scrutton, 1992, 1997; Fuller and Jenkins, 2007). *Flindersipora* is up to 70 cm tall (Fuller and Jenkins, 2007). *Moorowipora* and *Arrowipora*, cerioid forms with short septal spines and tabulae, resemble tabulates (Fuller and Jenkins, 1994, 1995; Sorauf and Savarese, 1995). Contemporaneous *Harklessia* (Botomian–Toyonian = ~512.5 Ma) lacks both septa and tabulae (Hicks, 2006). These domical coralomorphs declined at the end-early Cambrian, and only a few middle and late Cambrian examples are known (Debrenne and Reitner, 2001).

3.3.4. Chaetetids

In addition to coralomorphs, modular calcified skeletons occur in chaetetid sponges. In general, chaetetids have tabulae but poorly developed - or no - septa, whereas tabulate corals possess tabulae and may or may not have well-developed septa (Scrutton, 1979, 1997; West, 2012). Ediacaran *Namapoikia* has been compared with both chaetetids and coralomorphs (Wood et al., 2002). Botomian *Labyrinthus* (Kobluk, 1979) and *Rosellatana* (Kobluk, 1984), which attached to archaeocyaths, were regarded as coralomorphs by Debrenne and Reitner (2001). However, their mode of increase, similar to longitudinal fission, suggests affinities with chaetetids more than corals (Scrutton, 1997; Pratt et al., 2001). *Cambrophylum* (Fritz and Howell, 1955) from the lower Dresbachian Stage of North America (Guzhangian Age, ~499 Ma), 30 mm wide and 15 mm high, with septa-like processes within the ‘corallites’ (Scrutton, 1979; Jell, 1984; Kobluk, 1984) has also been compared with chaetetids (Scrutton, 1997). *Solenopora* (Riding, 2004), first known in the early Darriwilian (~465 Ma) (Klappa and James, 1980) and common in the late Darriwilian (Kröger et al., 2017) and early Late Ordovician (Opalinski and Harland, 1981), may be the first abundant rock-forming chaetetid.

3.3.5. *Pelmatozoans*

Pelmatozoans (stem-bearing echinoderms) appear in Cambrian Epoch 2 (Zamora et al., 2013; Peel, 2017a). The eocrinoid *Kinercystis* from early Epoch 2 (~519 Ma) is an early example (Zamora et al., 2013, fig. 13.3). Echinoderms declined in the Guzhangian–Paibian and then recovered in the mid–late Furongian (Zamora et al., 2013). Echinoderms attached to firm substrates in the mid–late Cambrian (Brett et al., 1983; Zamora et al., 2010, 2017), but within Cambrian reefs they mostly occur as scattered fragments (James and Kobluk, 1978; Pruss and Knoll, 2017) and may have been passive filter feeders restricted to low-velocity currents (Smith, 1990). Only locally did they form reef fabric, e.g., in the basal parts of some late Cambrian LSM reefs (Spencer, 1996; Johns et al., 2007). Nonetheless, together with bryozoans, their holdfasts created reef substrates in the late Tremadocian (~480 Ma) (Adachi et al., 2011, fig. 6), Darriwilian (Pratt, 1989) and Late Ordovician (e.g., *Hemicosmites*, Kröger et al., 2014).

3.4. Mid–late Cambrian (“reef gap”) late Age 4/end Age 10 (510–485 Ma): microbial-lithistid

Mid–late Cambrian reefs have long been known to be dominantly microbial (Fagerstrom, 1987; Rowland and Shapiro, 2002; Riding, 2006). Lithistid sponges had been recognized in the mid–late Cambrian, but were often considered too small, too simple, and too scarce to be effective reef-builders (Fagerstrom, 1987; Pratt et al., 2001; Rowland and Shapiro, 2002). Consequently, the interval between archaeocyath decline at the end of the early Cambrian and the appearance of large skeletal reefs in the mid-Ordovician came to be regarded as a major ‘reef gap’, the longest of the Phanerozoic (Copper, 2001; Stanley, 2001; Rowland and Shapiro, 2002; Kiessling, 2009; James and Wood, 2010). This view also tended to overlook the Early Ordovician reefs documented by Toomey (1970), Copper (1974), Toomey and Nitecki (1979) and others, that contain large domes and columns of *Pulchrellamina* as well as conspicuous conical lithistids (*Archaeoscyphia*) and calathiids (*Calathium*). Lithistid sponges as a whole constitute a long-ranging polyphyletic group of calcified demosponges (Pisera, 2002; Pisera and Lévi, 2002; Morrow and Cárdenas, 2015; Schuster et al., 2015) and continue to build reefs at the present-day (Maldonado et al., 2015). All known Cambrian lithistids, however, belong to the Family Anthaspidellidae, characterized by conical morphology and a distinctive ladderlike spicule structure. In overall shape and size, anthaspidellids broadly resemble archaeocyaths, and this similarity can extend to their reef-building role (Hong et al., 2016; Lee et al., 2016a). Both are commonly closely associated with, and volumetrically subordinate to, calcimicrobes and other microbial carbonates in reefs (Kruse and Zhuravlev, 2008; Kruse and Reitner, 2014; Adachi et al., 2015; Lee et al., 2016a).

For 25 Myr, throughout the mid–late Cambrian, skeletal reefs were dominated by cuplike lithistid sponges, always in association with microbial carbonates. The earliest known lithistid is *Rankenella* from near the Age 4/5 boundary (Kruse, 1983, 1996). A series of studies, beginning with Hamdi et al. (1995) (Lee et al., 2016a and references therein) (Fig. 1; Table 1), recognized lithistid contributions to mid–late Cambrian reefs. Lithistids were supported by their cemented basal attachments and interlocking spicules, and they could develop external ridged ornamentation that may have increased nutrient intake (Church, 2017). In Section 4 below, we summarize well-described mid–late Cambrian lithistid reefs, and in Fig. 2 show examples of lithistids. In addition to lithistids, mid–late Cambrian reefs also locally contain abundant siliceous sponges that have not yet been identified, but appear to resemble keratose sponges (cf. Luo and Reitner, 2014), and create maze-like “macerate” structures (Lee et al., 2014a, 2015a, 2016c; Coulson and Brand, 2016).

Most early Cambrian skeletal reef-builders declined in the mid–late Cambrian. Radiocyaths disappeared in late Epoch 2 (middle Toyonian) (Kruse et al., 2015). Archaeocyaths and coralomorphs declined

Table 1
Examples of mid–late Cambrian lithistid sponge-microbial reefs.

Location	Age	Environment	Reef description	References
Dedebulak Formation, Kyrgyzstan	Stage 5?	Unknown	Lithistid sponge <i>Orolinocyathus-Epiphyton</i> reef	Teslenko et al. (1983)
Zhangxia Fm., Shandong Province, China	Upper Stage 5-lower Guzhangian	Shallow subtidal	Lithistid sponge (<i>Rankenella</i>)-calcimicrobe (<i>Epiphyton</i>)-stem-group cnidarian (<i>Cambroctonus</i>) reef; meter-scale lens embedded within microbial-dominant reef; framework formed by mutual attachment of <i>Rankenella</i> and <i>Cambroctonus</i> with <i>Epiphyton</i> ; encrusted by microstromatolite and microbial clots; sponges occur between mesocots	Woo (2009); Adachi et al. (2015), Lee et al. (2016a)
Daegi Fm., Taebaeksan Basin, Korea	Drumian	Shallow subtidal inner platform	Thrombolite-sponge reef; sponges mostly <i>Anthaspidellidae</i> ; constructed primarily by <i>Epiphyton</i>	Hong et al. (2012, 2016)
Ranken Limestone, Georgina Basin, Australia	Upper Drumian	Bioclast-rich ramp debris	<i>Angusticellularia-Taninia</i> -lithistid sponge (<i>Rankenella</i>) reef; extent uncertain, possibly a few decimeters; sponges mainly passive reef builders within matrix formed by <i>Angusticellularia</i> and <i>Taninia</i>	Kruse and Reitner (2014)
Deh-Molla Formation, Iran	Lower Paibian	Subtidal, platform margin	Lithistid sponge (<i>Rankenella</i>)- <i>Girvanella</i> reef; sponges occupy up to 50% of reef volume and form framework by their mutual attachment; frameworks encrusted by <i>Girvanella</i> and/or microstromatolite	Hamdi et al. (1995), Kruse and Zhuravlev (2008)
Bonanza King Formation, Nevada and California, USA	Lower Paibian	Tidal channel within shallow subtidal ramp	Lithistid sponge (<i>Gallatinospongia</i>)-dendrolite reef; sponges in situ and enclosed by dendrolite; sponges occupy up to 30% in area	Shapiro and Rigby (2004)
Wilberns Formation, Texas, USA	Jiangshanian	Shallow marine	Lithistid sponge (<i>Wilberncyathus</i>)- <i>Girvanella</i> - <i>Taninia</i> meter-scale reef; minor <i>Taninia</i> , <i>Epiphyton</i> , <i>Renalcis</i> ; sponges constitute up to 30% of reef volume	Johns et al. (2007)
Notch Peak Formation, Nevada, USA	Lower Stage 10	Shallow subtidal	<i>Anthaspidellid</i> sponge-microbial reef; sponges form framework; pendent <i>Angusticellularia</i> grew beneath sponges; microstromatolites fill the interspaces between sponge frameworks and sometimes occur within spongocoels	Mrozek et al. (2003); Lee et al. (in prep.)
Dotsero Formation, Colorado, USA	Upper Stage 10	Shallow marine	<i>Girvanella</i> - <i>Taninia</i> -lithistid sponge (<i>Wilberncyathus</i>) reef; thin veneer over <i>Girvanella</i> stromatolite	Johns et al. (2007)

substantially (Zhuravlev et al., 1993; Scrutton, 1997), but both groups persisted. Two examples of archaeocyaths occur in the Drumian (Wood et al., 1992; ~500–504 Ma, Bassett-Butt, 2016) and late Cambrian (Furongian) (Debrenne et al., 1984) of Antarctica, but the group did not regain its former prominence or form reefs (Bambach et al., 2002) and is not known in the Ordovician. Mid-late Cambrian conical coralomorphs include *Cothonian* and *Lipopora* in Age 4 and *Tretocylichne* in Age 5 (Park et al., 2016, figs. 4, 5), and the similar genus *Cambroctoconus* in late Age 5/early Drumian (~504 Ma) (Park et al., 2011) which was a reef-building component (Lee et al., 2016a). *Cambrotrypa* (Fritz and Howell, 1959; Bolton and Copeland, 1963; Scrutton, 1979; Jell, 1984), from the middle Cambrian of North America, could be a recumbent cerioid to phaceloid coral (Scrutton, 1997). In contrast to archaeocyaths, coralomorphs (e.g., *Amsassia*) recovered in the Early Ordovician.

3.5. Early–Middle Ordovician (485–460 Ma): microbial-lithistid-calathiid-pulchrilaminid

Early Ordovician lithistid-microbial reefs broadly resemble those of the mid-late Cambrian, but the lithistids are larger (e.g., *Archaeoscyphia*) (Fig. 4) and an additional cuplike group, the calathiids, appeared, exemplified by *Calathium* from the late Tremadocian/early Floian (Church, 1974). The major change at this time, however, was the appearance of a variety of encrusting-domical forms that included pulchrilaminids, coralomorphs (*Amsassia*), and the first definite appearances of tabulate corals (*Lichenaria*), bryozoans (*Nekhorosheviella*) and stromatoporoids (*Cystostroma*). These encrusting skeletons progressively transformed reef structure (Kröger et al., 2017). Pulchrilaminids (*Pulchrilamina*, *Zondarella*) can be up to 1 m in size, half as large again as early Cambrian *Flindersipora*. These reefs locally formed skeletal frameworks and hosted cryptic benthic organisms (Hong et al., 2014, 2015; Li et al., 2017b) (Fig. 4). This significant increase in

skeletal reef-builders has been regarded as an initial phase of the GOBE, foreshadowing the subsequent importance of reef-building stromatoporoids and tabulate corals (Webby, 2002; Webby et al., 2004; Adachi et al., 2011).

3.5.1. Receptaculitids, radiocyaths, cyclocrinatids

These calcified cup-like, pear-shaped, or globular macrofossils differ in age, and their affinities (generally suggested to be sponge or algal) continue to be debated. Radiocyaths are restricted to the early Cambrian (Kruse et al., 2015), where they are generally minor reef components. They were initially compared with archaeocyaths (Bedford and Bedford, 1934), with which they are associated in reef construction; but they differ in structure (Debrenne et al., 1970). Receptaculitids appear ~478 Ma (Tremadocian/Floian) and *Calathium* is a common frame-builder in Early Ordovician reefs (Church, 1974, 2009; Toomey and Nitecki, 1979; Cañas and Carrera, 1993; Li et al., 2014, 2015; Shen and Neuweiler, 2018). Ordovician receptaculitids are diverse and have been subdivided into three main groups: ischaditids, receptaculitids, soanitids (or calathiids) (Nitecki et al., 2004, fig. 31.1). They are locally important in later Ordovician reefs (e.g., Kröger et al., 2014), and range to the late Paleozoic. Cyclocrinatids mainly occur in the mid-late Ordovician and were diverse (e.g., *Apidium*, *Coelosphæridium*, *Cyclocrinites*) near the Sandbian–Katian transition (Nitecki et al., 2004).

Receptaculitids, radiocyaths and cyclocrinatids have been compared because their walls are composed of numerous small plates that create a distinctively patterned external surface and bear internal rods. These structural similarities are greatest between radiocyaths and receptaculitids (Nitecki and Debrenne, 1979), and more superficial in cyclocrinatids (Beadle, 1988). Cyclocrinatids have long been compared with dasycladalean algae (Stolley, 1896), whereas receptaculitids (Austin, 1845; Billings, 1865) and radiocyaths (Bedford and Bedford, 1934) were both initially regarded as sponges. This long-established




Fig. 4. Summary of Cambrian and Early Ordovician reefs. (A) Size of reef-building sponges. (B) Size of reefs in log-scale. (C) Pie diagrams of main reef-building components (microbial vs. skeletal). See Table 1 for data sources.

distinction was challenged by the proposal that the receptaculitid *Ischadites* is - like cyclocrinids - a dasycladalean alga (Kesling and Graham, 1962; Byrnes, 1968). Support for this view not only resulted in it being applied to receptaculitids in general, but also in cyclocrinids being regarded as receptaculitids (Nitecki, 1970, 1972; Nitecki and Debenne, 1979; Rietschel and Nitecki, 1984).

Such sweeping interpretations attracted some opposition. Rietschel (1977) agreed that receptaculitids are algae but considered that they are not dasycladaleans. Beadle (1988) argued that receptaculitids and dasycladaleans are 'entirely unrelated'. These discussions were often complicated by being a part of the parallel debate about archaeocyath affinities (Rowland, 2001). Nitecki et al. (1999, 2004) suggested that cyclocrinids are algae similar to dasyclads, but that receptaculitids are neither algae nor sponges. This latter view was also often considered for archaeocyaths (Vologdin and Zhuravleva, 1947; Zhuravleva and Myagkova, 1987) before they became generally regarded as sponges (Debenne et al., 2015b).

One of the difficulties in determining receptaculitid affinities is the morphological heterogeneity of the group, ranging from enclosed pyriform to bowl-shaped bodies (Byrnes, 1968). Calathiids including *Calathium*/*Soanites* are clearly conical or cup-like (Myagkova, 1966; Church, 1991). This morphology, common in filter-feeding organisms (Billings, 1865; Myagkova, 1966; Li et al., 2015, 2017b), supports Billings' (1865) early view that *Calathium* is a sponge. A parsimonious view might be that radiocyaths (and cuplike receptaculitids, i.e., calathiids) are sponges, and that cyclocrinids (perhaps together with enclosed receptaculitids) are algae. If radiocyaths and receptaculitids are related, then radiocyath decline at the end of the early Cambrian, followed by the Early Ordovician appearance of the calathiids, could be a further example of a group that was reduced in diversity and abundance during the mid-late Cambrian but recovered in the Early Ordovician. More information is required to elucidate these possibilities.

3.5.2. Bryozoans

The early history of bryozoans is problematic. The affinity of Dresbachian (Fritz, 1948) (~499 Ma) *Archaeotrypa* (Fritz, 1947) is uncertain; it may be a bryozoan or echinoderm (see Scrutton, 1979; Jell, 1984; Kobluk, 1984; Rozanov and Zhuravlev, 1992; Zhuravlev et al., 1993). *Pywackia*, reported by Landing et al. (2010) from Cambrian Stage 10 of Mexico has been compared with octocorals (Taylor et al., 2013; but see Landing et al., 2015). The earliest currently unquestioned bryozoan is *Prophyllodictya* from lower Tremadocian wackestones of South China (Ma et al., 2015), and the earliest reef-building bryozoan is late Tremadocian (~479 Ma) laminar *Nekhorosheviella semisphaerica*, also from South China (Cuffey and Zhu, 2010; Adachi et al., 2012; Cuffey et al., 2013, fig. 2.4a). *Nekhorosheviella* mutually encrusts, can occur on lithistids, and together with crinoid holdfasts forms small mounds 30 cm wide and 10 cm high, similar to Middle Ordovician pelmatozoan patch reefs (Pratt, 1989; Adachi et al., 2011). Larger bryozoan reefs appear in the late Darriwilian (Pitcher, 1964).

3.5.3. Pulchrellaminids

Late Tremadocian (~480 Ma) *Pulchrellamina* (Toomey and Ham, 1967) and Dapingian *Zondarella* (Keller and Flügel, 1996) are by far the largest Early Ordovician skeletons, although they often enclose or are interlayered with sediment. Pulchrellaminids share similarities with stromatoporoids, but their characteristic features such as very thin latilaminae and 'more loosely aggregated meshwork of skeletal elements, including slender, upwardly tapering, spinose rods that are spiculelike' suggest they are an independent group (Webby, 2015).

3.5.4. Corals

Lichenaria, generally considered the earliest tabulate coral (Scrutton, 1997; Elias et al., 2008), is recorded from the early Tremadocian (Pratt and James, 1982) (their Green Head Mound occurs in the lower Tremadocian Watts Bight Formation. See Knight et al., 2008, p.

118). However, Laub (1984) regarded the earliest *Lichenaria* as Darriwilian, and therefore contemporaneous with other tabulates such as *Billingsaria* and *Eofletcheria* (Desrochers and James, 1989; Pickett and Percival, 2001; Kröger et al., 2017). The coralomorph *Amsassia* (Carrera et al., 2017) may have been confused with *Lichenaria*, but differs in growth mode and has been suggested to be an alga (Sun et al., 2014).

3.6. Late Middle–Late Ordovician (460–444 Ma): stromatoporeid-bryozoan-tabulate-receptaculitid-microbial

Stromatoporoids appear in the Early Ordovician (early Floian) as small specimens of the labechiid *Cystostroma* attached to *Calathium* (Li et al., 2017b). Much larger labechiids (*Labechia*, *Pachystylostroma*) and also clathrodictyiids (*Pseudostyloclyon*) appear in the late Darriwilian ~460 Ma (Pitcher, 1964). Together with bryozoans (*Batostoma*) and tabulate corals (*Billingsaria*), as well as the chaetoid *Solenopora*, they constructed large Darriwilian/Sandbian reefs, such as the classic Chazy examples in Vermont (Pitcher, 1964; Finks and Toomey, 1969; Kapp, 1975), and the upper Sandbian Carters Formation of Tennessee (Alberstadt et al., 1974). In addition to their size and abundance, mid-Ordovician stromatoporoids, bryozoans and tabulates are also notable for their laminar and domical shapes (Copper, 1974; Fagerstrom, 1987; Webby, 2002; Webby et al., 2004; Adachi et al., 2011) which transformed substrate colonization and vertical succession in reefs (Pitcher, 1964; Finks and Toomey, 1969; Alberstadt et al., 1974; Copper, 1974; Hong et al., 2017, 2018; Kröger et al., 2017). Rugose corals also appear in the late Darriwilian, but did not become common until the Sandbian–Katian (Baars et al., 2012). All these groups diversified further in the Late Ordovician, together with encrusting echinoderms, cuplike receptaculitids, and globular cyclocrinids. Skeletal reefs with bivalves, siliceous sponges, and the coralomorph *Amsassia*, also occur during this time interval (Lee et al., 2014c, 2016b; Sun et al., 2014). These developments were part of continued GOBE diversification in the Middle to Late Ordovician that resulted in complex and well-structured benthic communities (Droser and Sheehan, 1997; Webby, 2002; Harper, 2006). Lithistids, often in association with *Calathium* and microbial carbonates continued to be locally abundant in late Darriwilian reefs (Alberstadt et al., 1974; Klappa and James, 1980; Desrochers and James, 1989; Brunton and Dixon, 1994; Liu et al., 2003; Li et al., 2017c), but the volumetric importance of these components appears to have declined by the Late Ordovician as stromatoporoids, corals and bryozoans increased (Webby, 2002; Adachi et al., 2011).

4. Mid-late Cambrian lithistid sponge-microbial reefs

The nature and distribution of lithistid sponge-microbial reefs are central to our discussion of mid-late Cambrian environments and biotas. Following their discovery in lower Furongian strata of Iran (Hamdi et al., 1995), mid-late Cambrian LSM reefs have been recognized worldwide (Mrozek et al., 2003; Shapiro and Rigby, 2004; Johns et al., 2007; Kruse and Zhuravlev, 2008; Hong et al., 2012, 2016; Kruse and Reitner, 2014; Adachi et al., 2015; Lee et al., 2016a) (Table 1). In contrast to early Cambrian archaeocyath-microbial reefs and Early Ordovician lithistid sponge-microbial reefs, which have been studied in detail for more than 100 years, middle-late Cambrian LSM reefs remain much less well-known. Here we briefly summarize the composition and structure of four relatively well-described examples.

4.1. Zhangxia Formation (upper Stage 5, Cambrian Series 3), Shandong, China

The oldest LSM reefs currently known occur in the lower part of the Zhangxia Formation (Shandong Province, China) (Woo, 2009; Adachi et al., 2015) where they form lenses ~150 cm thick within microbial mounds (Lee et al., 2016a). Based on the trilobite biozone *Lioparia*, they belong to the upper part of Stage 5 of Cambrian Series 3. At this level,




Fig. 5. Examples of mid-late Cambrian reef-building sponges. (A) *Rankenella zhangxianensis* (upper Stage 5; Zhangxia Formation), Beiquanzi section, Shandong Province, China. (B) *R. mors* (upper Drumian; Ranken Limestone), locality NTGS4648, Georgina Basin, Australia (after Kruse and Reitner, 2014, fig. 5A). (C) *R. hamdii* (lower Paibian; Deh-Molla Formation), Shahmirzad section, northern Iran (after Kruse and Zhuravlev, 2008, fig. 7D). (D) Unidentified anthaspidellid sponge (Stage 10; Notch Peak Formation), Arrow Canyon section, Nevada, USA. Coin is 19 mm in diameter.

the Zhangxia Formation is mainly composed of bioturbated wackestone and oolitic, skeletal, and peloidal packstone-grainstone, deposited on a shallow subtidal carbonate platform (Woo, 2009; Lee et al., 2015b). Metazoans include the lithistid sponge *Rankenella zhangxianensis* Lee et al., 2016d (Family Anthaspidellidae, Order Orchocladina), the stem-group cnidarian *Cambroctoconus orientalis* Park et al., 2011, and unidentified siliceous sponges (Fig. 5A). Microbial components include *Epiphyton*, microstromatolite, and undifferentiated microbialite (Lee et al., 2016a) (Fig. 6A, B). *R. zhangxianensis* occupies up to 20% of the reef volume, is mostly conical in shape with smooth outlines, and is 3–21 mm in diameter and up to 610 mm in length. *C. orientalis* (8–11 mm in diameter, 11–13 mm in length; Park et al., 2011) is shaped like an octagonal cone and occupies < 3% of the reef volume. The unidentified siliceous sponges (~1% of reef volume) consist of irregularly shaped spicule networks resembling keratose sponges (cf. Luo and Reitner, 2014). Small dendritic calcified epiphytaceans are a major component, up to 20% of the reef volume. Microstromatolites are characterized by thin convex-up laminae of alternating dark and light gray micrite. Undifferentiated microbialite, composed of clotted-peloidal fabric with faint laminae, possibly represents poorly preserved microstromatolite. The main framework builders are the lithistids, *Epiphyton* and stem-group cnidarians (Lee et al., 2016a). The lithistids and stem-group cnidarians locally mutually attach (Fig. 5A), whereas *Epiphyton* is either mutually attached or attached to the sponges. These metazoan-calcimicrobial frameworks were encrusted and stabilized by microstromatolite (Fig. 6A, B), and probably created at least a few centimeters of syndepositional relief.

4.2. Ranken Limestone (upper Drumian, Cambrian Series 3), Georgina Basin, Australia

Reef fabrics formed by lithistid sponges and microbes in the Ranken Limestone are only known from loose blocks described by Kruse and Reitner (2014) who supposed that the individual reefs would not have exceeded a few decimeters in size. The Ranken Limestone consists of cross-bedded bioclast-oid-intraclast grainstone and subordinate bioclastic wackestone and appears to be a ramp deposit. Biostratigraphic control is poor, but relationships with adjacent deposits suggest a late Drumian age (Kruse and Reitner, 2014). The reef fabric mainly consists of microstromatolite and the anthaspidellid sponge *Rankenella mors* Kruse, 1983 (Kruse and Reitner, 2014) (Fig. 5B), which is mostly explanate in shape and only about 1 mm thick, although conical examples are also present. Domical or locally columnar masses of microstromatolite, up to 11 mm thick and 26 mm across, consist of light and dark laminae with patches of calcified microbes (*Angusticellularia*, *Taninia*). The Ranken reef is therefore mainly microbial and the role of *R. mors* appears to have been subordinate, mainly providing a substrate for stromatolite growth. Elsewhere in the Georgina Basin, *R. mors* inhabited low-energy anaerobic mudstone environments in the mid-Cambrian Arthur Creek Formation (Kruse, 1996; Debrenne and Reitner, 2001).

4.3. Deh-Molla Formation (lower Paibian, Furongian), northern Iran

These generally small LSM reefs, less than a few tens of centimeters




Fig. 6. Photomicrographs of lithistid sponges and microstromatolites in mid-late Cambrian reefs. (A, B) *Rankenella zhangxianensis*-*Epiphyton*-*Cambroctoconus orientalis* reef (upper Stage 5; Zhangxia Formation), Beiquanzi section, Shandong Province, China. (C, D) *Anthaspidellid* sponge-microbial reef (Stage 10; Notch Peak Formation), Arrow Canyon section, Nevada, USA.

in height and width (Hamdi et al., 1995; Kruse and Zhuravlev, 2008), occur in the Mila (renamed Deh-Molla) Formation (Geyer et al., 2014). The reef-bearing interval overlies the *Prochuangia* trilobite biozone, suggesting an early Furongian (Paibian) age (Peng et al., 1999). The platform margin on which it developed preserves microbial boundstone/floatstone with abundant demosponges (Geyer et al., 2014). The main reef components are *Rankenella hamdii* (anthaspidellid sponge) and *Girvanella* (calcified cyanobacterium). *R. hamdii* occupies up to 50% of the reef volume (Hamdi et al., 1995). It is conical, bowl or digitate in shape (Fig. 5C), 4–31 mm in diameter, and up to 80 mm in height and mutually attaches, forming frameworks that are commonly thickly encrusted by *Girvanella*, columnar microstromatolites and clotted-peloidal fabric. These encrustations mainly occur on the dermal surfaces of the sponges, but sometimes also on the gastral surfaces (Fig. 5C).

4.4. Notch Peak Formation (Stage 10, Furongian), Nevada, USA

Anthaspidellid sponge-microbialite reefs occur in the upper part of Notch Peak Formation (Mrozek et al., 2003). Oolitic and bioclastic packstone-grainstone and flat-pebble conglomerates, as well as stromatolites and the lithistid sponge-microbial reefs, are locally well-preserved within an otherwise dolomitized succession. The facies association suggests a shallow subtidal environment. The reefs are domical and a few meters thick, commonly occurring with columnar stromatolites (Mrozek et al., 2003). They contain the *Eoconodontus notchpeakensis* conodont fauna, suggesting a late Furongian (Age 10) age (Mrozek et al., 2003; Dattilo et al., 2004). Three main components of

these LSM reef are unidentified lithistid sponge, microstromatolite, and the calcified microbe *Angusticellularia* (Mrozek et al., 2003; J-H.L., personal observation) (Figs. 5D and 6C, D). The lithistids are mainly thin walled (~5 mm) upward widening cups (often 3 cm, but up to 14 cm, in diameter). They do not appear to closely resemble any described lithistid genus, but display typical anthaspidellid-type spicule networks (J-H.L., personal observation). The microstromatolites consist of thin alternating light-dark laminae, grading to faintly clotted/laminated fabric. Dark pendent *Angusticellularia* is common on the lower surfaces of the sponges. Keratose-like siliceous sponges occur rarely within the lithistid spongocoels. The main frame-builders of these Notch Peak reefs are lithistids, which appear to have been firmly mutually attached. The downward growing *Angusticellularia*, although forming only very thin crusts, suggests sediment-free space beneath the growing sponges. Bioturbated micrite, skeletal fragments and unidentified siliceous sponges fill the spongocoels. Microstromatolites occupy much of the space between sponges (Fig. 6C), and thin microstromatolite layers locally encrust sponge dermal surfaces within spongocoels (Fig. 6D).

4.5. Summary

Although much further research is required, in structure and composition these mid-late Cambrian LSM reefs quite closely resemble Early Ordovician examples built by microbes and the lithistid *Archaeoscyphia*. The above examples suggest several general characteristics of late Cambrian LSM reefs: 1. Decimeter- to meter-scale lensoid to domal shape. 2. Co-occurrence with, and often within,

microbial reefs. 3. Mutually attached frameworks formed by sponges. 4. Microstromatolites encrusting and stabilizing the sponge frameworks. 5. Formation in shallow subtidal packstone-grainstone environments.

5. Mid-late Cambrian dysoxia-hypoxia and sponge-microbial reefs

5.1. Evidence for low-oxygen conditions

A minimum level of oxygen required to maintain aerobic metabolism has long been linked to animal diversification during the Proterozoic–Cambrian transition (Berkner and Marshall, 1965; Rhoads and Morse, 1971; Knoll and Carroll, 1999; Canfield et al., 2007; Payne et al., 2009, 2011; Dahl et al., 2010; Maloof et al., 2010a; Knoll, 2011; Sperling et al., 2013; Ader et al., 2014; Mills and Canfield, 2014; Xiao, 2014; Zhang et al., 2014, 2018; Chen et al., 2015; Li et al., 2017a; Shields, 2017). It has been suggested that ocean oxygenation promoted animal diversification in the early Ediacaran ~630 Ma (Sahoo et al., 2012) and early-mid (541–509 Ma) Cambrian (Li et al., 2017a) and that ‘modern-like oxygen levels’ appeared by ~521 Myr ago (Chen et al., 2015). However, evidence from iron speciation (Sperling et al., 2015a), Mo-isotopes (Dahl et al., 2010) and alteration of deep sea basalts (Stolper and Keller, 2018) provide compelling evidence that persistent elevated oxygen levels are likely a later Paleozoic phenomenon. It seems that oxygenation of the oceans during the Ediacaran–Cambrian transition was likely a protracted and fluctuating series of oxic-anoxic/euxinic phases (Fike et al., 2006; Canfield et al., 2007, 2008; Shen et al., 2008; Jiang et al., 2009; Halverson et al., 2010; Wood et al., 2015; Zhang et al., 2018; Wood and Erwin, 2018). This secular pattern was further complicated by local variations in basinal and shelf anoxia (Goldberg et al., 2005; Schröder and Grotzinger, 2007; Canfield et al., 2008; Kendall et al., 2015; Sperling et al., 2015a; Jin et al., 2016; Och et al., 2016; Cheng et al., 2017). Moreover, the tendency toward low seawater oxygen concentrations subsequently appears to have increased during the Cambrian (Boyle et al., 2014; Cremonese et al., 2014), and persisted into the Ordovician (Railsback et al., 1990; Strauss, 2006; Kah and Bartley, 2011; Thompson and Kah, 2012; Wallace et al., 2017) (Fig. 1). Based on extensive black shale occurrences, Berry and Wilde (1978) inferred widespread and recurrent episodes of marine hypoxia during the mid-late Cambrian and Early Ordovician. They compared these conditions with those of the Cretaceous, attributed them to elevated temperatures, and suggested that they did not decline until cooling promoted mid-Ordovician ocean ventilation (Fig. 1). Black (or dark-colored) shales have been widely recognized in lower Paleozoic strata worldwide (e.g., Leggett, 1980; Wilde and Berry, 1984; Berry et al., 1986, 1989; Wilde et al., 1989; Fyfe and Pickerill, 1993; Saltzman et al., 2000; Egenhoff et al., 2015). The Chengjiang (ca. 520 Ma), Burgess Shale (ca. 508 Ma) and Fezouata (ca. 480 Ma) biotas, as well as other Burgess-Shale type biotas, which record a large proportion of early animal taxa, are preserved in organic-rich shales that are thought to reflect low-oxygen conditions (Conway Morris, 1986; Gaines et al., 2005; Gaines and Droser, 2010; Van Roy et al., 2010), even though other factors could have been necessary to preserve these fossils (Gaines, 2014). Some of these shales are significant sources of petroleum (e.g., Buchardt and Lewan, 1990; Mi et al., 2007; Ryder et al., 2014), and gas (e.g., Yu et al., 2009; Schovsbo et al., 2011; Huang et al., 2012; Pashin et al., 2012; Pool et al., 2012). Preservation of abundant organic matter in these shales suggests that low-oxygen conditions were prevalent throughout their deposition. Extinctions of archaeocyaths (Zhuravlev and Wood, 1996) and trilobites (Palmer, 1984; Westrop and Ludvigsen, 1987) in the Cambrian have been linked to anoxia.

This broad interpretation of a tendency to low oxygen marine conditions during the interval between the early Cambrian and mid-Ordovician has received support from studies of biogeochemically cycled redox-sensitive elements (C, N, S, U isotopic values; and Zn/Fe, Mo, Ce/Ce*, Th/U abundances), particularly for the mid-late Cambrian

(Perfetta et al., 1999; Saltzman et al., 2000; Gill et al., 2011; Dahl et al., 2014; Saltzman et al., 2015). Well documented examples of correspondence between black shale deposition, faunal change, and geochemical indicators of hypoxia include the ~510 Ma Sinsk and ~500 Ma SPICE events. Incursions of anoxic water onto shelves at ~510 Ma have been inferred from negative carbon isotope (Montañez et al., 2000; Guo et al., 2010) and positive $\delta^{34}\text{S}$ (Hough et al., 2006) values. Similar age $\delta^{15}\text{N}$ values, among the lowest of the Phanerozoic, may suggest low-oxygen photic zone conditions (Algeo et al., 2014, fig. 1). Interpretation of the ~500 Ma Steptoean Positive Carbon Isotope Excursion (SPICE) as reflecting low-oxygen conditions in shelf waters (Saltzman et al., 2000) is supported by coeval positive $\delta^{34}\text{S}$ sulfate excursions (Gill et al., 2007, 2011; Hurtgen et al., 2009), decreased Mo enrichment (Gill et al., 2011), and negative $\delta^{238}\text{U}$ excursion (Dahl et al., 2014). In addition, the Drumian (middle Cambrian) negative carbon isotope excursion (DICE) (~505 Ma) and associated Mo, U, and V values have been linked to euxinic conditions that extended into the photic zone (Pagès and Schmid, 2016). Cambrian–Ordovician boundary (~485 Ma) slope carbonates in Newfoundland have Ce/Ce* and Th/U ratios and N and U-isotope values consistent with dysoxic conditions (Azmy et al., 2014, 2015). The inference of Berry and Wilde (1978), that the tendency for shelf biotas to experience influxes of low-oxygen waters only diminished in the mid-Ordovician as ocean circulation improved, is supported by oxygen (Railsback et al., 1990) and sulfur (Kah et al., 2016) isotope studies as well as Th/U ratios in carbonates (Marenco et al., 2016), and has been linked to global cooling (Trotter et al., 2008) (Fig. 1).

These changes are widely considered to have been important in ushering in the GOBE (McKenzie et al., 2014; Saltzman et al., 2015). Edwards et al. (2017) suggest that atmospheric oxygen concentrations were low (~12%) during the Early Ordovician and doubled to about 25% by the Darriwilian, coinciding with increased biodiversity. Based on sea-surface temperature (Trotter et al., 2008, fig. 3) and modeled estimates of atmospheric oxygen (Edwards et al., 2017, fig. 2), we calculated the dissolved oxygen concentrations in seawater during the Ordovician (Fig. 7). These values are generally near 100 μM during the Early Ordovician, approximately one third of present-day seawater. They begin to rise ~465 Ma (early Darriwilian) and reach a peak at 451 Ma (early Katian), before declining steeply to 446 Ma (late Katian). This overall trend broadly corresponds with that of marine invertebrate diversity (Sepkoski Jr., 1995, fig. 1; Webby et al., 2004, fig. 1.1). Our calculated dissolved oxygen values (Fig. 7) from Edwards et al. (2017, fig. 2) are mainly for the Ordovician, but also indicate ~12% atmospheric oxygen in the late Cambrian. Based on the minimum requirements of animals (lower value) and pervasive subsurface anoxia (upper value), Sperling et al. (2015b, fig. 3) (also Stolper and Keller, 2018) estimated overall Cambrian atmospheric oxygen levels to have been even lower, in the range 15–40% of present atmospheric level (= 3–8% oxygen). These average values can be expected to have resulted in much lower levels of dissolved oxygen, seasonally and locally. Even in the present-day ocean, under 21% atmospheric O_2 , hypoxia commonly develops (Rabalais et al., 2010) and oxygen-deficient waters threaten coral reefs (Altieri et al., 2017). There is evidence for early to middle Cambrian oxygen-minimum zones (Guilbaud et al., 2018). It is therefore likely that localized and seasonal conditions resulted in fluctuations well below the average level of dissolved oxygen (~100–115 $\mu\text{M/L}$) that we calculated (Fig. 7) for the Early Ordovician, and which is already hypoxic according to the definition of Tyson and Pearson (1991).

Terms commonly used to describe dissolved oxygen levels include anoxic, suboxic, dysoxic/hypoxic, oxic/normoxic (e.g., Tyson and Pearson, 1991; Middleburg and Levin, 2009; Rabalais et al., 2010; Sperling et al., 2015a). These terms, the thresholds defining them, and the units used to measure them (e.g., mg/L, $\mu\text{M/kg}$, mL/L, $\mu\text{M/L}$) often differ according to author and research field. Tyson and Pearson (1991, table 2) pointed out that terms such as dysoxic, dysaerobic and hypoxic




Fig. 7. (A) Ordovician marine invertebrate diversity (Sepkoski Jr., 1995, fig. 1). (B) Dissolved oxygen concentrations in seawater calculated from sea-surface temperature (Trotter et al., 2008, fig. 3) and atmospheric oxygen (Edwards et al., 2017, fig. 2) data (see Supplementary Oxygen and Temperature Data).

had been applied to environment, biofacies and physiology respectively. Over time, however, some of these distinctions have become blurred. The following terms (for environments, facies, oxygenation levels) and thresholds were recommended by Tyson and Pearson (1991, table 2) in mL/L, at 10 °C, 1 atm pressure, with 1.029 density of seawater (conversions to other widely used units are shown in brackets):

oxic: 8.0–2.0 mL/L (= 12.03–3.01 mg/L, 376–94 μM /L, 365.4–91.35 μM /kg, 11.69–2.92 ppm)

dysoxic: 2.0–0.2 mL/L (= 3.01–0.30 mg/L, 94–9.4 μM /L, 91.35–9.14 μM /kg, 2.92–0.29 ppm)

suboxic: 0.2–0.0 mL/L (= 0.30–0 mg/L, 9.4–0 μM /L, 9.14–0 μM /kg, 0.29–0 ppm)

anoxic: ≥ 0.0 mL/L (= ≥ 0 in all cases)

Unlike reef-builders, mobile organisms can move to avoid localized environmental changes. Furongian and Early Ordovician reefs in Newfoundland, for example, occur in close association with skeletons of trilobites and cephalopods (see Pruss and Knoll, 2017). Several present-day groups, including shrimps and crabs, can detect and avoid oxygen deficient sea water (Renaud, 1986). Mobile nekton on the northern Gulf of Mexico continental shelf avoids oxygen levels < 2 mg/L (1.4 mL/L) (Pavela et al., 1983; Renaud, 1986), and fish avoidance generally occurs at oxygen concentrations 0.5 to 2 mg/L (0.35 to 1.4 mL/L) (Levin et al., 2009). Similarly, various motile organisms, including ctenophores, crustaceans, and cephalopods, can tolerate low oxygen levels (Ekau et al., 2010, table 2). Variability in oxygen concentrations can be more limiting than the absolute value (Matabos et al., 2012). There is evidence that late Cambrian groups such as olenids (Fortey, 2000) and

phosphatocopids (Williams et al., 2011) were dysoxia-tolerant. Large present-day arthropods, such as lobsters, are relatively tolerant of oxygen levels as low as 2 ppm, though 4 ppm is generally considered to be the minimum (Bayer et al., 1998). Our calculation of ~100 μM oxygen (3.2 ppm) in the Early Ordovician is approaching the upper limit of dysoxic (2.92 ppm) based on Tyson and Pearson (1991, table 2). This suggests that average oxygen levels in Early Ordovician seawater could have supported large arthropods but that seasonal and local variations could readily have resulted in oxygen limitation.

5.2. Factors promoting low-oxygen conditions in shallow marine environments

Interrelated factors generally contributing to shallow marine deoxygenation include global warming, increased marine productivity, thermal stratification, and expansion of shallow platform areas (Schmittner et al., 2008; Keeling et al., 2010; Praetorius et al., 2015). Global warmth, that characterized the mid-Cambrian to mid-Ordovician (Frakes et al., 1992, p. 14) and promoted low-oxygen conditions, has been linked to high sea-level and high CO_2 “greenhouse” conditions during commencement of the first Phanerozoic supercycle (Fischer, 1984; Landing, 2012b; Nance and Murphy, 2013). Modeled estimates suggest that CO_2 (Bernier, 2009) and temperature (Royer et al., 2004, fig. 4) peaked in the mid-Cambrian. Oxygen isotope thermometry indicates earliest Ordovician sea-surface temperatures (Shields et al., 2003; Kasting et al., 2006) in excess of 40 °C (Trotter et al., 2008). Earlier additional episodic warming due to CO_2 increase has been

attributed to extensive continental flood basalt volcanism, such as the late Cambrian Age 4 Kalkarindji Large Igneous Province in Australia (510.7 ± 0.6 Ma, Jourdan et al., 2014) (Glass and Phillips, 2006; Hough et al., 2006). As shown by Mesozoic Ocean Anoxic Events (OAEs), clear links can exist between anoxia and Large Igneous Provinces (LIPs). However, further studies are needed to identify Paleozoic LIPs (Percival et al., 2015, fig. 1) and disparities in timing between early Botomian anoxia (> 514 Ma; Peng et al., 2012), the Hawke's Bay regression (~ 511 Ma), and the Kalkarindji eruption (~ 510 Ma) recommend caution in linking these events.

Low oxygen areas develop or persist seasonally or continuously beneath upwelling regions, associated with the upper parts of oxygen minimum zones, and in warm oceans that are more stratified, hold less oxygen and experience greater advection of oxygen-poor source waters (Levin et al., 2009). In addition to lowering oxygen solubility, higher temperatures make water less dense, promoting thermal stratification that in turn reduces ventilation (Erbacher et al., 2001). Warming also favors biological productivity in the water column (e.g., phytoplankton blooms) which increases carbon burial, and thus the likelihood of bottom-water hypoxia (Pedersen and Calvert, 1990; Sen Gupta et al., 1996). Global Cambrian sea-level rise (Miller et al., 2005) promoted development of extensive, shallow epeiric carbonate platforms (Peters and Gaines, 2012), that tend to limit shallow-water circulation (Irwin, 1965) and expand the environments subject to thermal stratification (Allison and Wright, 2005; Allison and Wells, 2006). In Laurentia, mid-Cambrian sea-level rise established extensive platforms that locally persisted into the Late Ordovician (Bond et al., 1989; Morgan, 2012). This 'Great American Carbonate Bank' enclosed intrashelf basins (e.g., Markello and Read, 1981) prone to dysoxia-anoxia, and black mudstones accumulated on its slopes during eustatic highs (Landing, 2012a, 2012b; Lavoie et al., 2012; Miller et al., 2012; Peters and Gaines, 2012). These conditions ameliorated by the mid-Ordovician, as indicated by early Darriwilian sulfur isotope values consistent with ocean ventilation that appear to coincide with lower sea-surface temperatures (Kah et al., 2016), as well as with decrease in strontium isotope values (Edwards et al., 2015). Ordovician cooling and oxygenation, coincident with diversification of the "Paleozoic" and "Modern" faunas (Trotter et al., 2008), also marked the end of sponge-microbial reef dominance in the early Paleozoic (Brunton and Dixon, 1994). This trend continued until widespread oxygenation of the deep ocean occurred in the mid-Paleozoic (Dahl et al., 2010).

5.3. Low oxygen levels and lithistid sponges

Lithistids are a polyphyletic group of demosponges with interlocked siliceous spicules (desmas) that form a rigid skeleton (Hinde, 1883; Lévi, 1991; Debrenne and Reitner, 2001; Pisera, 2002; Pisera and Lévi, 2002; Hill et al., 2013). They are represented by about 46 extant genera (Maldonado et al., 2016). Present-day lithistids can dominate benthic faunas at depths of ~ 150 – 1800 m (Kelly, 2007), and locally aggregate into reefs (Conway et al., 1991) similar to ancient examples (Brunton and Dixon, 1994). Dense intertwined clusters of the lithistid *Leiodermatium pfeifferae*, with thin (3–9 mm) erect foliose branches up to ~ 80 cm high and 1 m wide, at depths of ~ 750 m in the western Mediterranean, have been compared with Mesozoic lithistid reefs (Maldonado et al., 2015, 2016). However, since the anthaspidellid lithistids that dominated Cambrian–Ordovician reefs went extinct in the Permian, caution is required when comparing lithistid reefs of different ages (Pisera, 2002).

Data concerning the oxygen requirements of present-day lithistids are limited; more information is available concerning other demosponges and also hexactinellids. Many sponges survive periods of anoxia. In experiments, demosponges not only tolerate hypoxia, but appear to have higher survival rates under hypoxic conditions than under normoxia (Gunda and Janapala, 2009). *Halichondria panicea*, for example, grows under oxygen in the range 0.5–4% of present atmospheric level

(Mills et al., 2014). Similar examples are documented from the wild. For example, in Saanich Inlet, western Canada, hexactinellids survive anoxia (oxygen concentrations < 0.2 mL/L) (Tunnicliffe, 1981), readily tolerate hypoxia (< 1.4 mL/L) (Chu and Tunnicliffe, 2015), and generally appear to do well in oxygen as low as 0.5 mL/L (i.e., severe hypoxia) (Jackson Chu, pers. comm, 2016). Thus, overall, at least some sponges dominate hypoxic systems and tolerate periodic anoxia (Hoffmann and Schläppy, 2011; Steckbauer et al., 2011; Riedel et al., 2014).

Sponge cells together with their associated symbiotic microbes often experience anaerobic conditions when pumping activity stops, and can maintain internal suboxic and locally hypoxic conditions (Hoffmann and Schläppy, 2011; Lavy et al., 2016). Sponges typically use only a small fraction of available oxygen for respiration (Hoffmann et al., 2005) and their ability to tolerate low oxygen levels may be linked to facultatively anaerobic mitochondria (Mentel et al., 2014). Some sponges, including lithistids (Brück et al., 2012) harbor large numbers of bacterial symbionts (Imhoff and Trüper, 1976; Fan et al., 2012) that can constitute up to 40% of sponge biomass (Taylor et al., 2007; Schmitt et al., 2012) and provide significant nutrition. These include anaerobic bacteria that occupy "anoxic niches" (Schläppy et al., 2010). The complex lithistid microbiome (Hentschel et al., 2012) is a source of pharmaceutical metabolites (Thomas et al., 2010).

Molds of sponges (Gehling and Rigby, 1996) and phosphatized sponges (Li et al., 1998; Yin et al., 2015) occur in the Ediacaran. Spicules that have been reported (Brasier et al., 1997) may have been misidentified (Antcliffe et al., 2014; Muscente et al., 2015). Biomarkers (Love et al., 2009; Sperling et al., 2010) suggest that demosponges and weakly calcified sponge-grade metazoans (Maloof et al., 2010b) appeared even earlier (Erwin et al., 2011), consistent with sponges having low oxygen requirements (Mills and Canfield, 2014). Oxygen increase during the Ediacaran–Cambrian transition (Fike et al., 2006; Chen et al., 2015) does not appear to have been sustained (see Section 5.1). The appearance of archaeocyaths in the early Cambrian, and their relatively abrupt decline coincident with the anoxic Sinsk Event (Zhuravlev and Wood, 1996) suggests that they may have been less tolerant than lithistids to low oxygen conditions. In contrast, lithistids appear unaffected by mid-late Cambrian and Early Ordovician extinction events attributed to anoxia (Saltzman et al., 2015). The only gradual increase observed in sponge size through the mid-Cambrian to early Ordovician (Fig. 3) could reflect the need to maintain the advantage of small body size under low-oxygen conditions (Payne et al., 2011).

Stromatoporoids and corals were important reef-builders from the mid-Ordovician to Late Devonian (Fagerstrom, 1987; Wood, 1999; James and Wood, 2010). The lithistid-microbial association reappeared sporadically; e.g., in the Late Ordovician (Brunton and Dixon, 1994; Leinfelder et al., 2002, p. 185), Late Silurian (de Freitas et al., 1999), and in Early Carboniferous reefs (e.g., Mundy, 1994; Webb, 1999) following Late Devonian stromatoporoid extinction. Similarly, a variety of sponges together with microbes occurs in reefs following the end-Permian extinction (Brayard et al., 2011). Probably the most conspicuous Mesozoic development of lithistid-microbial reefs was in the mid-late Jurassic, together with hexactinellid sponges (Leinfelder et al., 2002). Lithistid-microbial reefs are common in poorly oxygenated environments in the Upper Jurassic of Germany and Middle to Upper Jurassic of Spain and Portugal (Leinfelder et al., 1994, 1996) at depths of 30–90 m, coeval with shallower water coral-sponge-microbial reefs (Leinfelder, 2001, fig. 18). This was a period of relatively warm conditions (Jenkyns et al., 2012) when reduced oceanic circulation is inferred to have promoted black shale formations and dysaerobic reefs (Leinfelder et al., 1994, 1996). Lithistid-microbial reefs could therefore represent a default "low oxygen" Phanerozoic reef community.

Stromatoporoids may have had different oxygen requirements from lithistids. Present-day hypercalcified sponges (sclerosponges), considered to be relict stromatoporoids (Stearn, 2015b), preferentially

occur in shallow water (~20 m) with corals (Reitner, 1993; Quinn and Kojis, 1999). Together with light dependence in Paleozoic stromatoporoids (Kershaw, 2015; Stearn, 2015a) and the depth-dependent occurrence of hypercalcified sponges in Jurassic reefs (e.g., their dominance in shallow marine reefs and absence in deep, poorly oxygenated Jurassic lithistid-microbial reefs) (Fürsich and Werner, 1991; Werner et al., 1994), it can be postulated that stromatoporoids preferred shallow, oxygen-rich environments.

5.4. Mid-Cambrian to Early Ordovician reef development

It is now apparent that the mid-late Cambrian “reef gap” was occupied by LSM reefs, and that early Cambrian archaeocyath-microbial reefs and Early Ordovician LSM reefs were broadly similar in structure to those of the mid-late Cambrian. Cambrian to Early Ordovician reef development prompts a number of key questions: why did archaeocyaths decline; why were lithistids the only significant metazoan reef builders during the mid-late Cambrian; what factors could have limited other reef builders during this period; and what prompted the rise of new skeletal reef-builders in the Early Ordovician?

Archaeocyath decline has been linked to one or more of sea-level fall (Palmer and James, 1979), anoxia (Zhuravlev and Wood, 1996; Zhuravlev, 2001b), and ocean acidification (Knoll and Fischer, 2011); all of which have also tentatively been related to the Kalkarindji flood basalt eruptions (Glass and Phillips, 2006; Hough et al., 2006). The subsequent mid-late Cambrian interval, with reefs largely dominated by calcimicrobes and microbial carbonates, has been regarded as a post-extinction phase following archaeocyath demise, in which microbes represent a “disaster flora” (Fagerstrom, 1987; Zhuravlev, 1996; Wood, 1999). Although it has been speculated that the microbes may have outcompeted metazoan reef-builders during this interval (Riding and Zhuravlev, 1994; Zhuravlev, 1996; Rowland and Shapiro, 2002), this alone would not account for its ~25 Myr duration, which far exceeds recovery from much larger subsequent extinctions. For example, the end-Permian extinction event aftermath is not generally thought to have exceeded 5 Myr, and was significantly less in the case of metazoan-rich reefs which appeared within 1.5 Myr (Brayard et al., 2011).

In reviewing factors to explain the prolonged ‘resurgence of microbialites and the absence of reef-building metazoans’ during the mid-Cambrian to early Ordovician, Rowland and Shapiro (2002, p. 116–120) concluded that nutrient deficiency, elevated levels of CO₂ and temperature (including associated reduction in oxygen availability), and lower levels of Mg/Ca ratio in seawater could have been involved. Webby (2002) recognized that Ordovician low oxygen prolonged microbial carbonates and slowed metazoan diversification until the Late Ordovician. Increased bioturbation has been related to the return of widespread anoxic and sometimes euxinic conditions in the early-mid Paleozoic ocean (Boyle et al., 2014; Lenton et al., 2014), and Saltzman et al. (2015) noted that the Cambrian and Ordovician radiations are separated by persistent oceanic anoxia and elevated extinction rates. Pruss et al. (2010) linked the scarcity of calcified skeletons in Furongian–Tremadocian seas, in which subsurface water masses were ‘commonly dysoxic to anoxic’, to reduction in carbonate saturation. This interpretation of lowered saturation for CaCO₃ minerals in surface waters during episodes of anoxia stems from efforts to account for the scarcity of “robust skeletons” during the Early Triassic (Fischer et al., 2007; Knoll et al., 2007). Since anaerobic microbial processes can increase alkalinity, it has been suggested that times when deep anoxic seawater was geologically widespread could have had higher carbonate saturation (Ω) that promoted seafloor precipitation of CaCO₃ minerals, and that under these conditions of increased subsurface alkalinity ‘the Ω of overlying surface waters should be reduced’ (Higgins et al., 2009). Accordingly, Knoll and Fischer (2011, pp. 75, 78) suggested that the limited quantity of hypercalcified organisms (‘massively calcifying sponges and cnidarians, and calcareous algae’) and metazoan reefs in late Cambrian and earliest Ordovician oceans may, at least in part,

reflect depressed carbonate saturation in surface seawater. A difficulty with this interpretation is that such low carbonate saturation in shallow water should also have inhibited the formation of microbial carbonates and oolites, yet these deposits dominate many late Cambrian–early Ordovician successions, including those examined by Pruss et al. (2010).

Widespread and protracted hypoxia can critically affect the physiology of marine animals. We suggest that lithistids were more tolerant of low oxygen conditions than archaeocyaths, and also than most of the skeletal reef builders that flourished in the Middle to Late Ordovician, consistent with the mid-Cambrian to early Ordovician persistence of lithistid sponges. Subsequent invertebrate diversification and microbial carbonate decline during the Ordovician reflect the effects of progressive temperature reduction (Trotter et al., 2008; Rasmussen et al., 2016) and increased ocean ventilation (Young et al., 2016; Edwards et al., 2017) and oxygenation on metazoan diversification (Figs. 1, 3, 4, 7). Tolerance of hypoxia by lithistids and microbes, could therefore account for the LSM community's persistence during the greenhouse period of the early Paleozoic.

5.5. Overview

The pattern of early skeletal reef development outlined here commenced with *Cloudina* ~550 Ma in the late Ediacaran and continued until large stromatoporoid-coral reefs appeared ~465 Ma in the Darriwilian. We recognize six reef intervals during this time (Fig. 8). I: Late Ediacaran (~550–541 Ma) reefs contain *Cloudina* up to 15 cm in size, *Namacalathus* up to 3.5 cm and *Namapoikia* (although apparently limited to fissures) up to 1 m in extent. These organisms have not been recognized in the Paleozoic. II: So far as is known, reefs of the earliest Cambrian (Fortunian–mid-Age 2) lack skeletal metazoans except for “*Ladatheca*” worms. III: Archaeocyath sponges, from mid-Age 2 to within Age 4, commonly up to 15 cm in size, locally constructed relatively large frameworks (Fig. 4). IV: Late Age 4 to Age 10 lithistid sponges and the reefs they constructed were both initially relatively small (Figs. 4, 5), but increased in size in the late Cambrian (Age 10). V: Additions to lithistids in Early Ordovician reefs include cup-like calathids, bryozoans, *Amsassia*, *Lichenaria*, *Cystostroma* and *Pulchrilamina*. With the possible exception of *Namapoikia*, the latter four may represent the earliest massive laminar-domical skeletal reef-builders. Thus, from mid-late Cambrian to early Ordovician, lithistid-microbial reefs, appear to show progressive increase (Figs. 3, 4) in sponge size and reef size, as well as in diversity with the addition of non-lithistid reef-builders. VI: This interval commenced in the late Darriwilian with the appearance of stromatoporoid-coral-bryozoan reefs (Webby, 2002).

These various Ediacaran–Ordovician reefs are all potentially frame-reefs in the sense of Riding (2002). Their metazoan skeletons are tubular, coniform, globular, domical, and laminar in outline, and all appear to have been capable, at least partially, of mutual attachment (Pitcher, 1964; Rowland, 1984; Riding and Zhuravlev, 1995; Penny et al., 2014; Hong et al., 2015; Li et al., 2015; Lee et al., 2016a). Nonetheless, vertical-oriented forms (tubular, coniform, globular) require either mutual support or matrix (including microbial carbonate) support, whereas laminar to domical forms are more able to create their own stability (Kröger et al., 2017). Thus, Ediacaran–Ordovician reef building skeletons shift from dependence on matrix and/or framework support (erect tubular, globular, coniform) prior to the Early Ordovician, to self-stabilizing, encrusting laminar-domical forms (tabulate corals, bryozoans, stromatoporoids) in the mid-Ordovician.

Numerous areas for investigation arise from this overview of early Phanerozoic skeletal reef-building organisms. Much remains to be discovered concerning the affinities, ecology and history of Ediacaran organisms such as *Cloudina*, *Namacalathus* and *Namapoikia*. These appear to have suffered end-Ediacaran extinction (Amthor et al., 2003), but the apparent scarcity of reef-building animals in the earliest Cambrian requires confirmation. Similar questions of affinity and history




Fig. 8. Size patterns of late Ediacaran to Middle Ordovician reef-building skeletal organisms based on examples in Supplementary Table 1. Three major morphotypes are emphasized: tubular, coniform-globular, domical-laminar. Tubular *Cloudina* and “*Ladatheca*” occur in the late Ediacaran and early Cambrian, respectively. Coniform sponges (archaeocyaths, lithistids, *Calathium*), augmented by globular radiocyaths, are locally conspicuous in Cambrian and Early Ordovician reefs. Domical-laminar forms are represented by coralomorphs in Cambrian Age 2–4, and by pulchrlaminids, stromatoporoids, bryozoans and corals in the Ordovician. These data suggest peaks in skeletal reef-builder size in Reef Interval III and in Reef Intervals V and VI, with smaller skeletons during Reef Interval IV. Reef intervals: I, microbial-*Cloudina*; II, microbial-“*Ladatheca*”; III, microbial-archaeocyath; IV, microbial-lithistid; V, microbial-lithistid-calathiid-pulchrlaminid; VI, stromatoporoid-bryozoan-tabulate-receptaculitid-microbial.

surround Cambrian coralomorphs, and the early history of lithistids is also unclear. The earliest recorded lithistid, *Rankenella* from near the early–middle Cambrian boundary (Kruse, 1996), does not appear to be reefal. The earliest known reef-building lithistid is *Rankenella zhangxianensis* in the upper part of Stage 5 of Cambrian Series 3 (Lee et al., 2016a). Our focus on the mid–late Cambrian draws attention to environmental changes associated with archaeocyath decline in late Epoch 2, as well as those linked to reef diversification in the Tremadocian. It also directs attention toward lithistid reef abundance within the last 25 Myrs of the Cambrian. Most of the examples we emphasize here have been discovered or described since 2003 (Table 1). Certainly, many more remain to be discovered and will provide new insights into this neglected group and their reef-building abilities. Similarly, recent progress in understanding coralomorphs, bryozoans, tabulates, pulchrlaminids and stromatoporoids is important in assessing how laminar/domical morphotypes contributed to and changed reefs during the Ordovician. Commonly overlooked groups, such as radiocyaths, receptaculitids and cyclocrinids, are locally probably more significant reef components than is generally realized and deserve more attention. In this review we have often mentioned, but not emphasized, the significant role of microbial carbonates as major reef components associated with these skeletal organisms throughout the late Ediacaran to

mid-Ordovician. Competitive and mutualistic relationships between these closely associated but very different components provide fertile ground for research to elucidate substrate evolution and reef-building strategies at the dawn of the Phanerozoic. Not least, our emphasis of the significance of marine oxygenation, and its spatial and temporal patterns during this interval, raises numerous unresolved and tempting questions at the interfaces of geochemistry, geobiology and sedimentology.

6. Conclusions

Cambrian reefs were dominated by microbial carbonates (stromatolites, dendrolites, thrombolites), together with either calcareous (archaeocyath) or siliceous (lithistid) sponges capable of constructing interconnected skeletal frameworks. Heavily calcified archaeocyaths are relatively conspicuous in early Cambrian reefs. Following their decline, microbial carbonates continued to dominate reefs and the mid–late Cambrian was regarded as a significant gap in metazoan reef development. However, it is now recognized that lithistid sponges are widespread in mid–late Cambrian reefs. Although easy to overlook and often subordinate to microbial carbonates, lithistids built frameworks similar to archaeocyaths. This lithistid-microbial reef consortium




Fig. 9. Schematic patterns of Ediacaran–Ordovician skeletal reef structure and marine oxygenation based on data summarized here (see Figs. 1, 2, 7). Microbial-archaeocyath reefs include domical coralomorphs. Marine oxygenation is inferred to have stimulated reef diversification into and including the early Cambrian, whereas reduced oxygen availability in mid-late Cambrian reef environments is reflected by development of lithistid-microbial reefs. This continued, with addition of calathiids and pulchrlaminids, into the early Ordovician. Increased marine oxygenation in the mid-late Ordovician stimulated further and significant diversification of reef organisms, setting the scene for much of the Silurian and Devonian.

continued into the Early Ordovician, when lithistids became more diverse and reef-building was significantly augmented by bryozoans, tabulate corals, and sponges such as the archaeocyath-like *Calathium* and stromatoporoid-like *Pulchrlamina*. This Early Ordovician reef association can be seen as a transition from Cambrian lithistid-microbial reefs to the bryozoan-tabulate coral-stromatoporoid reefs that flourished in the late Darriwilian, ~460 Ma. Subsequently, these relatively large and heavily calcified skeletons, many of which are laminar to domical in overall morphology, continued to dominate reef-building for ~100 Myr, until the Late Devonian.

Geochemical evidence suggests that rise in marine oxygenation during the Ediacaran–Cambrian transition was not maintained, and was followed during the mid-late Cambrian by a widespread tendency to marine hypoxia-dysoxia. This supports the long-established inference that black shale intervals, linked to episodes of Cambrian trilobite extinction, reflect widespread anoxia. It is also consistent with the view that mid-late Cambrian “greenhouse” conditions, with temperatures increased by elevated CO_2 , led to marine stratification that reduced oxygen availability. Extensive carbonate platforms that developed in response to increased CO_2 , temperature and sea-level, are likely to have further restricted shallow-marine circulation in the mid-late Cambrian and Early Ordovician.

Many present-day silicified sponges, including lithistids, are well-adapted to low oxygen conditions, as are microbes; whereas corals and bryozoans generally are not. The mid-late Cambrian development and persistence of lithistid-microbial reefs could therefore reflect a prolonged tendency to low oxygen conditions in shallow marine environments. Oxygen limitation may have caused the decline of late early Cambrian archaeocyaths and also hindered their subsequent recovery. It is likely to have delayed the diversification of reef builders such as bryozoans and corals. The tendency toward marine hypoxia continued for at least 25 Myr, until the end of the Cambrian. Its gradual amelioration in the Early Ordovician can be linked to decline in sea-surface temperatures that promoted ocean ventilation. The addition of bryozoans, tabulates and stromatoporoids to lithistid-microbial reefs in the Early Ordovician reflects progressive marine oxygenation. Lithistid-microbial reefs did not disappear, but their period of early Paleozoic dominance was at an end by the late Darriwilian. Nonetheless, nearly

300 Myr later, during the Middle to Late Jurassic, a similar lithistid-microbial reef community reappeared in low-oxygen marine environments.

Marine oxygenation likely stimulated Ediacaran–Cambrian diversification, but “greenhouse” conditions subsequently reduced oxygen availability in shallow reefal environments, delaying mid-late Cambrian metazoan diversification, and prolonging development of lithistid-microbial reefs (Fig. 9). Marine oxygen level could therefore have been a key factor that initially facilitated early Cambrian diversification, and then – as oxygenation declined – restrained its subsequent advance until the Ordovician. This two-step tempo, paralleling that of coeval animal evolution in general, could account for the prolonged pause between the Cambrian Explosion and the GOBE.

Acknowledgements

We thank Jackson Chu and Henry Reiswig for advice regarding sponge ecology, Keun-Hyung Choi for information concerning arthropod oxygen requirements, and Liyuan Liang for calculating the dissolved oxygen values in Fig. 7 together with the Supplemental Data. We are indebted to three reviewers for valuable comments and suggestions, and to André Strasser for expert editorial guidance and encouragement. JHL's research was supported by National Research Foundation of Korea (grant 2016R1C1B1012104). This study is a contribution to IGCP Project 653 ‘The onset of the Great Ordovician Biodiversity Event’.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.earscirev.2018.04.003>.

References

- Adachi, N., Ezaki, Y., Liu, J., 2011. Early Ordovician shift in reef construction from microbial to metazoan reefs. *PALAIOS* 26, 106–114. <http://dx.doi.org/10.2110/palo.2010.p10-097r>.
- Adachi, N., Ezaki, Y., Liu, J., 2012. The oldest bryozoan reefs: a unique Early Ordovician skeletal framework construction. *Lethaia* 45, 14–23. <http://dx.doi.org/10.1111/j.>

- 1502-3931.2011.00268.x.
- Adachi, N., Kotani, A., Ezaki, Y., Liu, J., 2015. Cambrian Series 3 lithistid sponge-microbial reefs in Shandong Province, North China: reef development after the disappearance of archaeocyaths. *Lethaia* 48, 405–416. <http://dx.doi.org/10.1111/let.12118>.
- Ader, M., Sansjofre, P., Halverson, G.P., Busigny, V., Trindade, R.I.F., Kunzmann, M., Nogueira, A.C.R., 2014. Ocean redox structure across the Late Neoproterozoic Oxygenation Event: a nitrogen isotope perspective. *Earth Planet. Sci. Lett.* 396, 1–13. <http://dx.doi.org/10.1016/j.epsl.2014.03.042>.
- Adrain, J.M., McAdams, N.E.B., Westrop, S.R., 2009. Trilobite biostratigraphy and revised bases of the Tulean and Blackhillsian stages of the Ibexian Series, Lower Ordovician, western United States. *Memoirs of the Association of Australasian Palaeontologists* 37, 541–610.
- Alberstadt, L.P., Walker, K.R., Zurawski, R.P., 1974. Patch reefs in the Carters Limestone (Middle Ordovician) in Tennessee, and vertical zonation in Ordovician reefs. *Geol. Soc. Am. Bull.* 85, 1171–1182. [http://dx.doi.org/10.1130/0016-7606\(1974\)85<1171:PRITCL>2.0.CO;2](http://dx.doi.org/10.1130/0016-7606(1974)85<1171:PRITCL>2.0.CO;2).
- Algeo, T.J., Meyers, P.A., Robinson, R.S., Rowe, H., Jiang, G.Q., 2014. Icehouse-greenhouse variations in marine denitrification. *Biogeosciences* 11, 1273–1295. <http://dx.doi.org/10.5194/bg-11-1273-2014>.
- Allison, P.A., Wells, M.R., 2006. Circulation in large ancient epicontinental seas: what was different and why? *PALAIOS* 21, 513–515. <http://dx.doi.org/10.2110/palo.2006.506>.
- Allison, P.A., Wright, V.P., 2005. Switching off the carbonate factory: A-tidality, stratification and brackish wedges in epeiric seas. *Sediment. Geol.* 179, 175–184. <http://dx.doi.org/10.1016/j.sedgeo.2005.05.004>.
- Altieri, A.H., Harrison, S.B., Seemann, J., Collin, R., Diaz, R.J., Knowlton, N., 2017. Tropical dead zones and mass mortalities on coral reefs. *Proc. Natl. Acad. Sci.* 114, 3660–3665. <http://dx.doi.org/10.1073/pnas.1621517114>.
- Amthor, J.E., Grotzinger, J.P., Schröder, S., Bowring, S.A., Ramezani, J., Martin, M.W., Matter, A., 2003. Extinction of *Cloudina* and *Namacalathus* at the Precambrian-Cambrian boundary in Oman. *Geology* 31, 431–434. [http://dx.doi.org/10.1130/0091-7613\(2003\)031<0431:EOCANA>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(2003)031<0431:EOCANA>2.0.CO;2).
- Antcliffe, J.B., Callow, R.H., Brasier, M.D., 2014. Giving the early fossil record of sponges a squeeze. *Biol. Rev.* 89, 972–1004. <http://dx.doi.org/10.1111/brv.12090>.
- Austin, T., 1845. Note on Mr. Bowerbank's paper on the genus *Dunstervillea* (Bowerbank), with remarks on the *Ischadites Königii*, the *Tentaculites* and the *Conularia*. *Ann. Mag. Nat. Hist.* 15, 406–407.
- Azmy, K., Stouge, S., Brand, U., Bagnoli, G., Ripperdan, R., 2014. High-resolution chemostratigraphy of the Cambrian–Ordovician GSSP: enhanced global correlation tool. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 409, 135–144. <http://dx.doi.org/10.1016/j.palaeo.2014.05.010>.
- Azmy, K., Kendall, B., Brand, U., Stouge, S., Gordon, G.W., 2015. Redox conditions across the Cambrian–Ordovician boundary: elemental and isotopic signatures retained in the GSSP carbonates. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 440, 440–454. <http://dx.doi.org/10.1016/j.palaeo.2015.09.014>.
- Baars, C., Ghobadi Pour, M., Atwood, R.C., 2012. The earliest rugose coral. *Geol. Mag.* 150, 371–380. <http://dx.doi.org/10.1017/s0016756812000829>.
- Babcock, L.E., Peng, S.-C., Brett, C.E., Zhu, M.-Y., Ahlberg, P., Bevis, M., Robison, R.A., 2015. Global climate, sea level cycles, and biotic events in the Cambrian Period. *Palaeoworld* 24, 5–15. <http://dx.doi.org/10.1016/j.palwor.2015.03.005>.
- Bambach, R.K., 2006. Phanerozoic biodiversity mass extinctions. *Annu. Rev. Earth Planet. Sci.* 34, 127–155. <http://dx.doi.org/10.1146/annurev.earth.33.092203.122654>.
- Bambach, R.K., Knoll, A.H., Sepkoski Jr., J.J., 2002. Anatomical and ecological constraints on Phanerozoic animal diversity in the marine realm. *Proc. Natl. Acad. Sci.* 99, 6854–6859. <http://dx.doi.org/10.1073/pnas.0921509999>.
- Bambach, R.K., Knoll, A.H., Wang, S.C., 2004. Origination, extinction, and mass depletions of marine diversity. *Paleobiology* 30, 522–542. [http://dx.doi.org/10.1666/0094-8373\(2004\)030<0522:OEAMDO>2.0.CO;2](http://dx.doi.org/10.1666/0094-8373(2004)030<0522:OEAMDO>2.0.CO;2).
- Bassett-Butt, L., 2016. Systematics, biostratigraphy and biogeography of brachiopods and other fossils from the middle Cambrian Nelson Limestone, Antarctica. *GFF* 138, 377–392. <http://dx.doi.org/10.1080/11035897.2015.1094510>.
- Bayer, R., Riley, J., Donahue, D., 1998. The effect of dissolved oxygen level on the weight gain and shell hardness of new-shell American lobster *Homarus americanus*. *J. World Aquacult. Soc.* 29, 491–493. <http://dx.doi.org/10.1111/j.1749-7345.1998.tb00674.x>.
- Beadle, S.C., 1988. Dasyclads, cyclocrinids and receptaculitids: comparative morphology and paleoecology. *Lethaia* 21, 1–12. <http://dx.doi.org/10.1111/j.1502-3931.1988.tb01745.x>.
- Bedford, R., Bedford, W.R., 1934. New species of Archaeocyathinae and other organisms from the Lower Cambrian of Beltana, South Australia. In: *Memoirs of the Kyancutta Museum*. 1. pp. 1–7.
- Bengston, S., 2004. Early skeletal fossils. In: *Paleontological Society Papers*. 10. pp. 67–78. <http://dx.doi.org/10.1017/S1089332600002345>.
- Berkner, L.V., Marshall, L.C., 1965. On the origin and rise of oxygen concentration in the Earth's atmosphere. *J. Atmos. Sci.* 22, 225–261. [http://dx.doi.org/10.1175/1520-0469\(1965\)022<0225:OTOARO>2.0.CO;2](http://dx.doi.org/10.1175/1520-0469(1965)022<0225:OTOARO>2.0.CO;2).
- Berner, R.A., 1990. Atmospheric carbon dioxide levels over Phanerozoic time. *Science* 249, 1382–1386. <http://dx.doi.org/10.1126/science.249.4975.1382>.
- Berner, R.A., 2009. Phanerozoic atmospheric oxygen: new results using the GEOCARBSULF model. *Am. J. Sci.* 309, 603–606. <http://dx.doi.org/10.2475/07.2009.03>.
- Berry, W.B.N., Wilde, P., 1978. Progressive ventilation of the oceans—an explanation for the distribution of the Lower Paleozoic black shales. *Am. J. Sci.* 278, 257–275.
- Berry, W.B.N., Wilde, P., Quinby-Hunt, M.S., Orth, C.J., 1986. Trace element signatures in Dictyonema Shales and their geochemical and stratigraphic significance. *Nor. Geol. Tidsskr.* 66, 45–51.
- Berry, W.B.N., Wilde, P., Quinby-Hunt, M.S., 1989. Paleozoic (Cambrian through Devonian) anoxic biotopes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 74, 3–13. [http://dx.doi.org/10.1016/0031-0182\(89\)90016-3](http://dx.doi.org/10.1016/0031-0182(89)90016-3).
- Billings, E., 1865. Palaeozoic fossils. *Geol. Surv. Canada* 208–211.
- Bolton, T.E., Copeland, M.J., 1963. *Cambrotrypa* and *Bradoria* from the Middle Cambrian of western Canada. *J. Paleontol.* 1069–1070.
- Bond, G.C., Komzin, M.A., Steckler, M.S., Grotzinger, J.P., 1989. Role of thermal subsidence, flexure, and eustasy in the evolution of Early Paleozoic passive-margin carbonate platforms. In: Crevello, P.D., Wilson, J.L., Sarg, J.F., Read, J.F. (Eds.), *Controls on Carbonate Platform and Basin Development*. SEPM Special Publication 44, Tulsa, pp. 39–61.
- Bowring, S.A., Grotzinger, J.P., Isachsen, C.E., Knoll, A.H., Pelechaty, S.M., Kolosov, P., 1993. Calibrating rates of Early Cambrian evolution. *Science* 261, 1293–1298. <http://dx.doi.org/10.1126/science.11539488>.
- Boyle, R.A., Dahl, T.W., Dale, A.W., Shields-Zhou, G.A., Zhu, M., Brasier, M.D., Canfield, D.E., Lenton, T.M., 2014. Stabilization of the coupled oxygen and phosphorus cycles by the evolution of bioturbation. *Nat. Geosci.* 7, 671–676. <http://dx.doi.org/10.1038/ngeo2213>.
- Brasier, M., Green, O., Shields, G., 1997. Ediacarian sponge spicule clusters from south-western Mongolia and the origins of the Cambrian fauna. *Geology* 25, 303–306. [http://dx.doi.org/10.1130/0091-7613\(1997\)025<0303:esscfs>2.3.co;2](http://dx.doi.org/10.1130/0091-7613(1997)025<0303:esscfs>2.3.co;2).
- Brayard, A., Vennin, E., Olivier, N., Bylund, K.G., Jenks, J., Stephen, D.A., Bucher, H., Hofmann, R., Goudemand, N., Escarguel, G., 2011. Transient metazoan reefs in the aftermath of the end-Permian mass extinction. *Nat. Geosci.* 4, 694–697. <http://dx.doi.org/10.1038/ngeo1264>.
- Brett, C.E., Liddell, W.D., Derstler, K.L., 1983. Late Cambrian hard substrate communities from Montana/Wyoming: the oldest known hardground encrusters. *Lethaia* 16, 281–289. <http://dx.doi.org/10.1111/j.1502-3931.1983.tb02010.x>.
- Brück, W.M., Reed, J.K., McCarthy, P.J., 2012. The bacterial community of the lithistid sponge *Discodermia* spp. as determined by cultivation and culture-independent methods. *Mar. Biotechnol.* 14, 762–773. <http://dx.doi.org/10.1007/s10126-012-9443-6>.
- Brunton, F.R., Dixon, O.A., 1994. Siliceous sponge-microbe biotic associations and their recurrence through the Phanerozoic as reef mound constructors. *PALAIOS* 9, 370–387. <http://dx.doi.org/10.2307/3515056>.
- Buchardt, B., Lewan, M.D., 1990. Reflectance of vitrinite-like macerals as a thermal maturity index for Cambrian-Ordovician Alum Shale, southern Scandinavia. *AAPG Bull.* 74, 394–406.
- Byrnes, J.G., 1968. Notes on the nature and environmental significance of the Receptaculitaceae. *Lethaia* 1, 368–381.
- Cañas, F., Carrera, M., 1993. Early Ordovician microbial-sponge-receptaculitid bioherms of the Precordillera, western Argentina. *Facies* 29, 169–178. <http://dx.doi.org/10.1007/BF02536927>.
- Canfield, D.E., Poulton, S.W., Narbonne, G.M., 2007. Late-Neoproterozoic deep-ocean oxygenation and the rise of animal life. *Science* 315, 92–95. <http://dx.doi.org/10.1126/science.1135013>.
- Canfield, D.E., Poulton, S.W., Knoll, A.H., Narbonne, G.M., Ross, G., Goldberg, T., Strauss, H., 2008. Ferruginous conditions dominated later Neoproterozoic deep-water chemistry. *Science* 321, 949–952. <http://dx.doi.org/10.1126/science.1154499>.
- Carrera, M.G., Astini, R.A., Gomez, F.J., 2017. A lowermost Ordovician tabulate-like corallomorph from the Precordillera of western Argentina: a main component of a reef-framework consortium. *J. Paleontol.* 91, 73–85. <http://dx.doi.org/10.1017/jpa.2016.145>.
- Chen, X., Ling, H.F., Vance, D., Shields-Zhou, G.A., Zhu, M., Poulton, S.W., Och, L.M., Jiang, S.Y., Li, D., Cremonese, L., Archer, C., 2015. Rise to modern levels of ocean oxygenation coincided with the Cambrian radiation of animals. *Nat. Commun.* 6, 7142. <http://dx.doi.org/10.1038/ncomms8142>.
- Cheng, M., Li, C., Zhou, L., Feng, L., Algeo, T.J., Zhang, F., Romaniello, S., Jin, C., Ling, H., Jiang, S., 2017. Transient deep-water oxygenation in the early Cambrian Nanhua Basin, South China. *Geochim. Cosmochim. Acta* 210, 42–58. <http://dx.doi.org/10.1016/j.gca.2017.04.032>.
- Chu, J.W., Tunncliffe, V., 2015. Oxygen limitations on marine animal distributions and the collapse of epibenthic community structure during shoaling hypoxia. *Glob. Chang. Biol.* 21, 2989–3004. <http://dx.doi.org/10.1111/gcb.12898>.
- Church, S.B., 1974. Lower Ordovician patch reefs in western Utah. *Brigham Young Univ. Geol. Stud.* 21, 41–62.
- Church, S.B., 1991. A new Lower Ordovician species of *Calathium*, and skeletal structures of western Utah calathids. *J. Paleontol.* 65, 602–610. <http://dx.doi.org/10.1017/S0022366000030699>.
- Church, S.B., 2009. Problematic receptaculitid fossils from western Utah and eastern Nevada. In: Tripp, B.T., Krahulec, K., Jordan, J.L. (Eds.), *Geology and Geologic Resources and Issues of Western Utah*. Utah Geological Association Publication Vol. 38. pp. 55–66.
- Church, S.B., 2017. Efficient ornamentation in Ordovician anthaspidellid sponges. *Paleontol. Contrib.* 18, 1–8.
- Cloud, P.E., 1948. Some problems and patterns of evolution exemplified by fossil invertebrates. *Evolution* 2, 322–350. <http://dx.doi.org/10.1111/j.1558-5646.1948.tb02750.x>.
- Cohen, K.M., Finney, S.C., Gibbard, P.L., Fan, J.-X., 2013. The ICS international chronostratigraphic chart. *Episodes* 36, 199–204.
- Conway Morris, S., 1986. The community structure of the Middle Cambrian Phylloporid Bed (Burgess Shale). *Palaeontology* 29, 423–467.
- Conway Morris, S., 2006. Darwin's dilemma: the realities of the Cambrian 'explosion'. *Philos. Trans. R. Soc. Lon. B Biol. Sci.* 361, 1069–1083. <http://dx.doi.org/10.1098/rstb.2006.1846>.
- Conway Morris, S., Mattes, B.W., Chen, M., 1990. The early skeletal organism *Cloudina*: new occurrences from Oman and possibly China. *Am. J. Sci.* 290-A, 245–260.
- Conway, K.W., Barrie, J.V., Austin, W.C., Luternauer, J.L., 1991. Holocene sponge bioherms on the western Canadian continental shelf. *Cont. Shelf Res.* 11, 771–790. [http://dx.doi.org/10.1016/0278-4343\(91\)90079-L](http://dx.doi.org/10.1016/0278-4343(91)90079-L).
- Copper, P., 1974. Structure and development of early Paleozoic reefs. In: *Proceedings of the Second International Coral Reef Symposium*, pp. 365–386.
- Copper, P., 2001. Evolution, radiations, and extinctions in Proterozoic to mid-Paleozoic

- reefs. In: Stanley, G.D. (Ed.), *The History and Sedimentology of Ancient Reef Systems*. Kluwer Academic/Plenum Publishers, New York, pp. 89–119. http://dx.doi.org/10.1007/978-1-4615-1219-6_3.
- Coulson, K.P., Brand, L.R., 2016. Lithistid sponge-microbial reef-building communities construct laminated, upper Cambrian (Furongian) 'stromatolites'. *PALAIOS* 31, 358–370. <http://dx.doi.org/10.2110/palo.2016.029>.
- Cremonese, L., Shields-Zhou, G.A., Struck, U., Ling, H.-F., Och, L.M., 2014. Nitrogen and organic carbon isotope stratigraphy of the Yangtze platform during the Ediacaran–Cambrian transition in South China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 398, 165–186. <http://dx.doi.org/10.1016/j.palaeo.2013.12.016>.
- Cuffey, R.J., Zhu, Z., 2010. Yichang (Central China)–the oldest-known bryozoan reefs (mid-Early Ordovician), and comparison with the next-oldest (Garden Island, New York/Vermont, early Middle Ordovician). *Geol. Soc. Am. Abstr. Programs* 42, 64.
- Cuffey, R.J., Chuantao, X., Zhu, Z., Spjeldnaes, N., Hu, Z.-X., 2013. The world's oldest-known bryozoan reefs: late Tremadocian, mid-Early Ordovician; Yichang, Central China. In: Ernst, A., Schäfer, P., Scholz, J. (Eds.), *Bryozoan Studies 2010*. Springer-Verlag, Berlin/Heidelberg, pp. 13–27. http://dx.doi.org/10.1007/978-3-642-16411-8_2.
- Dahl, T.W., Hammarlund, E.U., Anbar, A.D., Bond, D.P., Gill, B.C., Gordon, G.W., Knoll, A.H., Nielsen, A.T., Schovsbo, N.H., Canfield, D.E., 2010. Devonian rise in atmospheric oxygen correlated to the radiations of terrestrial plants and large predatory fish. *Proc. Natl. Acad. Sci.* 107, 17911–17915. <http://dx.doi.org/10.1073/pnas.1011287107>.
- Dahl, T.W., Boyle, R.A., Canfield, D.E., Connelly, J.N., Gill, B.C., Lenton, T.M., Bizzarro, M., 2014. Uranium isotopes distinguish two geochemically distinct stages during the later Cambrian SPICE event. *Earth Planet. Sci. Lett.* 401, 313–326. <http://dx.doi.org/10.1016/j.epsl.2014.05.043>.
- Dattilo, B.F., Hlohowskyj, S., Ripperdan, R.L., Miller, J.F., Shapiro, R.S., 2004. Stratigraphic setting of an Upper Cambrian metazoan reef between the Nopah Formation to Goodwin Formation transition in southern Nevada. *Geol. Soc. Am. Abstr. Programs* 36, 368.
- de Freitas, T.A., Trettin, H.P., Dixon, O.A., Mallamo, M., 1999. Silurian System of the Canadian Arctic Archipelago. *Bull. Can. Petrol. Geol.* 47, 136–193.
- Debrenne, F., Reitner, J., 2001. Sponges, cnidarians, and ctenophores. In: Zhuravlev, A.Y., Riding, R. (Eds.), *Ecology of the Cambrian Radiation*. Columbia University Press, New York, pp. 301–325.
- Debrenne, F., Termier, H., Termier, G., 1970. Radiocyatha. Une nouvelle classe d'organismes primitifs du Cambrien inférieur. *Bulletin de la Société Géologique de France* 12, 120–125 (in French).
- Debrenne, F., Rozanov, A.Y., Webers, G.F., 1984. Upper Cambrian Archaeocyatha from Antarctica. *Geol. Mag.* 121, 291–299. <http://dx.doi.org/10.1017/S0016756800029186>.
- Debrenne, F.M., Gangloff, R.A., Lafuste, J.G., 1987. *Tabulaconus* Handfield: microstructure and its implication in the taxonomy of primitive corals. *J. Paleontol.* 61, 1–9. <http://dx.doi.org/10.1017/S0022336000028122>.
- Debrenne, F., Zhuravlev, A.Y., Kruse, P.D., 2015a. Archaeocyatha and Cribricyatha *nomen nudum*; taxa not Archaeocyatha, Radiocyatha, or Cribricyatha. In: Stearn, C.W. (Ed.), *Treatise on Invertebrate Paleontology, Part E, Porifera, Revised, Hypercalcified Porifera, Volumes 4 and 5*. The University of Kansas Paleontological Institute, Lawrence, Kansas, pp. 1105–1106.
- Debrenne, F., Zhuravlev, A.Y., Kruse, P.D., 2015b. General Features of the Archaeocyatha. In: Stearn, C.W. (Ed.), *Treatise on Invertebrate Paleontology, Part E, Porifera, Revised, Hypercalcified Porifera, Volumes 4 and 5*. The University of Kansas Paleontological Institute, Lawrence, Kansas, pp. 845–922.
- Desrochers, A., James, N.P., 1989. Middle Ordovician (Chazyan) bioherms and biostromes of the Mingan Islands, Quebec. In: Geldsetzer, H.H.J., James, N.P., Tebbutt, G.E. (Eds.), *Reefs: Canada and adjacent areas*. Canadian Society of Petroleum Geologists Memoir 13. Canadian Society of Petroleum Geologists, pp. 183–191.
- Drosdova, N.A., 1980. Algae in the Lower Cambrian organogenetic buildups of western Mongolia. In: *Transactions of the Joint Soviet-Mongolian Paleontological Expedition*. 10. pp. 1–140 (in Russian).
- Droser, M.L., Sheehan, P.M., 1997. Palaeoecology of the Ordovician radiation; resolution of large-scale patterns with individual clade histories, palaeogeography and environments. *Geobios* 20, 221–229. [http://dx.doi.org/10.1016/S0016-6995\(97\)80027-7](http://dx.doi.org/10.1016/S0016-6995(97)80027-7).
- Edwards, C.T., Saltzman, M.R., Leslie, S.A., Bergström, S.M., Sedlacek, A.R.C., Howard, A., Bauer, J.A., Sweet, W.C., Young, S.A., 2015. Strontium isotope ($^{87}\text{Sr}/^{86}\text{Sr}$) stratigraphy of Ordovician bulk carbonate: implications for preservation of primary seawater values. *Geol. Soc. Am. Bull.* 127, 1275–1289. <http://dx.doi.org/10.1130/b31149.1>.
- Edwards, C.T., Saltzman, M.R., Royer, D.L., Fike, D.A., 2017. Oxygenation as a driver of the Great Ordovician Biodiversification Event. *Nat. Geosci.* 10, 925–929. <http://dx.doi.org/10.1038/s41561-017-0006-3>.
- Egenhoff, S.O., Fishman, N.S., Ahlberg, P., Maletz, J., Jackson, A., Kolte, K., Lowers, H., Mackie, J., Newby, W., Petrowsky, M., 2015. Sedimentology of SPICE (Stephanian positive carbon isotope excursion): a high-resolution trace fossil and microfabric analysis of the middle to late Cambrian Alum Shale Formation, southern Sweden. In: Larsen, D., Egenhoff, S.O., Fishman, N.S. (Eds.), *Paying Attention to Mudrocks: Priceless! Geological Society of America Special Paper 515*. Geological Society of America, pp. 87–102. <http://dx.doi.org/10.1130/2015.2515.05>.
- Ekau, W., Auel, H., Pörtner, H.O., Gilbert, D., 2010. Impacts of hypoxia on the structure and processes in pelagic communities (zooplankton, macro-invertebrates and fish). *Biogeosciences* 7, 1669–1699. <http://dx.doi.org/10.5194/bg-7-1669-2010>.
- Elias, R.J., Lee, D.-J., Woo, S.-K., 2008. Corallite increase and mural pores in *Lichenaria* (Tabulata, Ordovician). *J. Paleontol.* 82, 377–390. <http://dx.doi.org/10.1666/06-114.1>.
- Erbacher, J., Huber, B.T., Norris, R.D., Markey, M., 2001. Increased thermohaline stratification as a possible cause for an ocean anoxic event in the Cretaceous period. *Nature* 409, 325–327. <http://dx.doi.org/10.1038/35053041>.
- Erwin, D.H., Laflamme, M., Tweedt, S., Sperling, E.A., Pisani, D., Peterson, K.J., 2011. The Cambrian conundrum: early divergence and later ecological success in the early history of animals. *Science* 334, 1091–1097. <http://dx.doi.org/10.1126/science.1206375>.
- Fagerstrom, J.A., 1987. *The Evolution of Reef Communities*. John Wiley and Sons, New York (600 pp).
- Faggetter, L.E., Wignall, P.B., Pruss, S.B., Newton, R.J., Sun, Y., Crowley, S.F., 2017. Trilobite extinctions, facies changes and the ROECE carbon isotope excursion at the Cambrian series 2–3 boundary, Great Basin, western USA. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 478, 53–66. <http://dx.doi.org/10.1016/j.palaeo.2017.04.009>.
- Fan, L., Reynolds, D., Liu, M., Stark, M., Kjelleberg, S., Webster, N.S., Thomas, T., 2012. Functional equivalence and evolutionary convergence in complex communities of microbial sponge symbionts. *Proc. Natl. Acad. Sci.* 109, E1878–E1887. <http://dx.doi.org/10.1073/pnas.1203287109>.
- Fike, D.A., Grotzinger, J.P., Pratt, L.M., Summons, R.E., 2006. Oxidation of the Ediacaran ocean. *Nature* 444, 744–747. <http://dx.doi.org/10.1038/nature05345>.
- Finks, R.M., Toomey, D.F., 1969. The paleoecology of Chazyan (lower Middle Ordovician) "reefs" or "mounds". State University College of Arts and Sciences, Plattsburg, New York State Geological Association Field Guide Book, pp. 93–120.
- Fischer, A.G., 1984. The two Phanerozoic supercycles. In: Berggren, W.A., Van Couvering, J.A. (Eds.), *Catastrophes and Earth History: The New Uniformitarianism*. Princeton University Press, Princeton, NJ, pp. 129–150.
- Fischer, W.W., Higgins, J.A., Pruss, S.B., 2007. Delayed biotic recovery from the Permian–Triassic extinction may have been influenced by a redox-driven reorganization of the marine carbonate system. *Geol. Soc. Am. Abstr. Programs* 39, 420.
- Fortey, R., 2000. Olenid trilobites: the oldest known chemoautotrophic symbionts? *Proc. Natl. Acad. Sci.* 97, 6574–6578. <http://dx.doi.org/10.1073/pnas.97.12.6574>.
- Frakes, L.A., Francis, J.E., Syktus, J.I., 1992. *Climate modes of the Phanerozoic*. Cambridge University Press (274 pp).
- Fritz, M.A., 1947. Cambrian Bryozoa. *J. Paleontol.* 21, 434–435.
- Fritz, M.A., 1948. Cambrian Bryozoa more precisely dated. *J. Paleontol.* 22, 373.
- Fritz, M.A., Howell, B.F., 1955. An Upper Cambrian coral from Montana. *J. Paleontol.* 29, 181–183.
- Fritz, M.A., Howell, B.F., 1959. *Cambrotrypa montanensis*, a Middle Cambrian fossil of possible coral affinities. *Proc. Geol. Assoc. Canada* 11, 89–93.
- Fuller, M.K., Jenkins, R.J.F., 1994. *Moorowipora chamberensis*, a coral from the Early Cambrian Moorowie Formation, Flinders Ranges, South Australia. *Trans. R. Soc. S. Aust.* 118, 227–235.
- Fuller, M.K., Jenkins, R.J.F., 1995. *Arrowipora fromensis* a new genus of tabulate-like coral from the Early Cambrian Moorowie Formation, Flinders Ranges, South Australia. *Trans. R. Soc. S. Aust.* 119, 75–82.
- Fuller, M., Jenkins, R., 2007. Reef corals from the Lower Cambrian of the Flinders Ranges, South Australia. *Paleaentology* 50, 961–980. <http://dx.doi.org/10.1111/j.1475-4983.2007.00682.x>.
- Fürsich, F.T., Werner, W., 1991. Palaeoecology of coralline sponge-coral meadows from the Upper Jurassic of Portugal. *Paläontol. Z.* 65, 35–69. <http://dx.doi.org/10.1007/BF02985773>.
- Fyffe, L.R., Pickerill, R.K., 1993. Geochemistry of Upper Cambrian–Lower Ordovician black shale along a northeastern Appalachian transect. *Geol. Soc. Am. Bull.* 105, 897–910. [http://dx.doi.org/10.1130/0016-7606\(1993\)105<0897:GOUCLO>2.3.CO;2](http://dx.doi.org/10.1130/0016-7606(1993)105<0897:GOUCLO>2.3.CO;2).
- Gaines, R.R., 2014. Burgess Shale-type preservation and its distribution in space and time. In: Laflamme, M., Schiffbauer, J.D., Darroch, S.A.F. (Eds.), *Reading and Writing of the Fossil Record: Preservation Pathways to Exceptional Fossilization. The Paleontological Society Papers 20*. The Paleontological Society, pp. 123–146.
- Gaines, R.R., Droser, M.L., 2010. The paleoredox setting of Burgess Shale-type deposits. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 297, 649–661. <http://dx.doi.org/10.1016/j.palaeo.2010.09.014>.
- Gaines, R.R., Kennedy, M.J., Droser, M.L., 2005. A new hypothesis for organic preservation of Burgess Shale taxa in the middle Cambrian Wheeler Formation, House Range, Utah. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 220, 193–205. <http://dx.doi.org/10.1016/j.palaeo.2004.07.034>.
- Gandin, A., Debrenne, F., 2010. Distribution of the archaeocyath-calcimicrobial bioconstructions on the Early Cambrian shelves. *Palaeoworld* 19, 222–241. <http://dx.doi.org/10.1016/j.palwor.2010.09.010>.
- Gehling, J.G., Rigby, J.K., 1996. Long expected sponges from the Neoproterozoic Ediacara fauna of South Australia. *J. Paleontol.* 70, 185–195. <http://dx.doi.org/10.1017/S0022336000023283>.
- Gerhardt, A.M., Gill, B.C., 2016. Elucidating the relationship between the later Cambrian end-Marjuman extinctions and SPICE event. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 461, 362–373. <http://dx.doi.org/10.1016/j.palaeo.2016.08.031>.
- Germes, G.J.B., 1972. New shelly fossils from Nama Group, South West Africa. *Am. J. Sci.* 272, 752–761. <http://dx.doi.org/10.2475/ajs.272.8.752>.
- Geyer, G., Bayet-Goll, A., Wilmsen, M., Mahboubi, A., Moussavi-Harami, R., 2014. Lithostratigraphic revision of the middle Cambrian (Series 3) and upper Cambrian (Furongian) in northern and Central Iran. *Newsl. Stratigr.* 47, 21–59. <http://dx.doi.org/10.1127/0078-0421/2014/0039>.
- Gill, B.C., Lyons, T.W., Saltzman, M.R., 2007. Parallel, high-resolution carbon and sulfur isotope records of the evolving Paleozoic marine sulfur reservoir. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 256, 156–173. <http://dx.doi.org/10.1016/j.palaeo.2007.02.030>.
- Gill, B.C., Lyons, T.W., Young, S.A., Kump, L.R., Knoll, A.H., Saltzman, M.R., 2011. Geochemical evidence for widespread euxinia in the later Cambrian ocean. *Nature* 469, 80–83. <http://dx.doi.org/10.1038/nature09700>.
- Glass, L.M., Phillips, D., 2006. The Kalkarindji continental flood basalt province: a new Cambrian large igneous province in Australia with possible links to faunal extinctions. *Geology* 34, 461–464. <http://dx.doi.org/10.1130/g22122.1>.
- Goldberg, T., Poulton, S., Strauss, H., 2005. Sulphur and oxygen isotope signatures of late Neoproterozoic to early Cambrian sulphate, Yangtze Platform, China: diagenetic

- constraints and seawater evolution. *Precambrian Res.* 137, 223–241. <http://dx.doi.org/10.1016/j.precamres.2005.03.003>.
- Grotzinger, J.P., Watters, W.A., Knoll, A.H., 2000. Calcified metazoans in thrombolite-stromatolite reefs of the terminal Proterozoic Nama Group, Namibia. *Paleobiology* 26, 334–359. [http://dx.doi.org/10.1666/0094-8373\(2000\)026<0334:CMITSR>2.0.CO;2](http://dx.doi.org/10.1666/0094-8373(2000)026<0334:CMITSR>2.0.CO;2).
- Guilbaud, R., Slater, B.J., Poulton, S.W., Harvey, T.H.P., Brooks, J.J., Nettersheim, B.J., Butterfield, N.J., 2018. Oxygen minimum zones in the early Cambrian Ocean. *Geochem. Persp. Lett.* 33–38. <http://dx.doi.org/10.7185/geochemlet.1806>.
- Gunda, V.G., Janapala, V.R., 2009. Effects of dissolved oxygen levels on survival and growth in vitro of *Haliclona pigmentifera* (Demospongiae). *Cell Tissue Res.* 337, 527–535. <http://dx.doi.org/10.1007/s00441-009-0843-5>.
- Guo, Q., Strauss, H., Liu, C., Zhao, Y., Yang, X., Peng, J., Yang, H., 2010. A negative carbon isotope excursion defines the boundary from Cambrian Series 2 to Cambrian Series 3 on the Yangtze Platform, South China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 285, 143–151. <http://dx.doi.org/10.1016/j.palaeo.2009.11.005>.
- Halverson, G.P., Wade, B.P., Hurtgen, M.T., Barovich, K.M., 2010. Neoproterozoic chemostratigraphy. *Precambrian Res.* 182, 337–350. <http://dx.doi.org/10.1016/j.precamres.2010.04.007>.
- Hamdi, B., Rozanov, A.Y., Zhuravlev, A.Y., 1995. Latest Middle Cambrian metazoan reef from northern Iran. *Geol. Mag.* 132, 367–373. <http://dx.doi.org/10.1017/S0016756800021439>.
- Harper, D.A.T., 2006. The Ordovician biodiversification: setting an agenda for marine life. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 232, 148–166. <http://dx.doi.org/10.1016/j.palaeo.2005.07.010>.
- Hentschel, U., Piel, J., Degnan, S.M., Taylor, M.W., 2012. Genomic insights into the marine sponge microbiome. *Nat. Rev. Microbiol.* 10, 641–654. <http://dx.doi.org/10.1038/nrmicro2839>.
- Hicks, M., 2006. A new genus of early Cambrian coral in Esmeralda County, southwestern Nevada. *J. Paleontol.* 80, 609–615. [http://dx.doi.org/10.1666/0022-3360\(2006\)80\[609:ANGOCJ\]2.0.CO;2](http://dx.doi.org/10.1666/0022-3360(2006)80[609:ANGOCJ]2.0.CO;2).
- Higgins, J.A., Fischer, W.W., Schrag, D.P., 2009. Oxygenation of the ocean and sediments: consequences for the seafloor carbonate factory. *Earth Planet. Sci. Lett.* 284, 25–33. <http://dx.doi.org/10.1016/j.epsl.2009.03.039>.
- Hill, D., 1964. The phylum Archaeocyatha. *Biol. Rev.* 39, 232–258. <http://dx.doi.org/10.1111/j.1469-185X.1964.tb00956.x>.
- Hill, M.S., Hill, A.L., Lopez, J., Peterson, K.J., Pomponi, S., Diaz, M.C., Thacker, R.W., Adamska, M., Boury-Esnault, N., Cardenas, P., Chaves-Fonnegra, A., Danka, E., De Laine, B.O., Formica, D., Hajdu, E., Lobo-Hajdu, G., Klontz, S., Morrow, C.C., Patel, J., Picton, B., Pisani, D., Pohlmann, D., Redmond, N.E., Reed, J., Richey, S., Riesgo, A., Rubin, E., Russell, Z., Rutzler, K., Sperling, E.A., di Stefano, M., Tarver, J.E., Collins, A.G., 2013. Reconstruction of family-level phylogenetic relationships within Demospongiae (Porifera) using nuclear encoded housekeeping genes. *PLoS One* 8, e50437. <http://dx.doi.org/10.1371/journal.pone.0050437>.
- Hinde, G.J., 1883. *Catalogue of the Fossil Sponges in the Geological Department of the British Museum (Natural History): With Descriptions of New and Little-Known Species.* Taylor and Francis, London (237 pp).
- Hoffmann, F., Schlöppy, M.-L., 2011. Sponges (Porifera) and sponge microbes. In: Thiel, V. (Ed.), *J. Reitner. Springer, Encyclopedia of Geobiology*, pp. 840–847. http://dx.doi.org/10.1007/978-1-4020-9212-1_194.
- Hoffmann, F., Larsen, O., Thiel, V., Rapp, H.T., Pape, T., Michaelis, W., Reitner, J., 2005. An anaerobic world in sponges. *Geomicrobiol. J.* 22, 1–10. <http://dx.doi.org/10.1080/01490450509022505>.
- Hofmann, H.J., Mountjoy, E.W., 2001. *Namacalathus-Cloudina* assemblage in Neoproterozoic Miette Group (Byng Formation), British Columbia: Canada's oldest shelly fossils. *Geology* 29, 1091–1094. [http://dx.doi.org/10.1130/0091-7613\(2001\)029\(1091:NCANIM\)2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(2001)029(1091:NCANIM)2.0.CO;2).
- Hong, J., Cho, S.-H., Choh, S.-J., Woo, J., Lee, D.-J., 2012. Middle Cambrian siliceous sponge-calcimicrobe buildups (Daegi Formation, Korea): metazoan buildup constituents in the aftermath of the Early Cambrian extinction event. *Sediment. Geol.* 253–254, 47–57. <http://dx.doi.org/10.1016/j.sedgeo.2012.01.011>.
- Hong, J., Choh, S.-J., Lee, D.-J., 2014. Tales from the crypt: early adaptation of crypto-biotic sessile metazoans. *PALAIOS* 29, 95–100. <http://dx.doi.org/10.2110/palo.2014.076>.
- Hong, J., Choh, S.-J., Lee, D.-J., 2015. Untangling intricate microbial-sponge frameworks: the contributions of sponges to Early Ordovician reefs. *Sediment. Geol.* 318, 75–84. <http://dx.doi.org/10.1016/j.sedgeo.2015.01.003>.
- Hong, J., Lee, J.-H., Choh, S.-J., Lee, D.-J., 2016. Cambrian Series 3 carbonate platform of Korea dominated by microbial-sponge reefs. *Sediment. Geol.* 341, 58–69. <http://dx.doi.org/10.1016/j.sedgeo.2016.04.012>.
- Hong, J., Choh, S.-J., Park, J., Lee, D.-J., 2017. Construction of the earliest stromatoporeoid framework: labechiid reefs from the Middle Ordovician of Korea. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 470, 54–62. <http://dx.doi.org/10.1016/j.palaeo.2017.01.017>.
- Hong, J., Oh, J.-R., Lee, J.-H., Choh, S.-J., Lee, D.-J., 2018. The earliest evolutionary link of metazoan bioconstruction: laminar stromatoporeoid-bryozoan reefs from the Middle Ordovician of Korea. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 492, 126–133. <http://dx.doi.org/10.1016/j.palaeo.2017.12.018>.
- Hough, M.L., Shields, G.A., Evins, L.Z., Strauss, H., Henderson, R.A., Mackenzie, S., 2006. A major sulphur isotope event at c. 510 Ma: a possible anoxia-extinction-volcanism connection during the Early-Middle Cambrian transition? *Terra Nova* 18, 257–263. <http://dx.doi.org/10.1111/j.1365-3121.2006.00687.x>.
- Huang, J., Zou, C., Li, J., Dong, D., Wang, S., Wang, S., Cheng, K., 2012. Shale gas generation and potential of the Lower Cambrian Qiongzhusi Formation in the southern Sichuan Basin, China. *Pet. Explor. Dev.* 39, 75–81. [http://dx.doi.org/10.1016/s1876-3804\(12\)60017-2](http://dx.doi.org/10.1016/s1876-3804(12)60017-2).
- Hurtgen, M.T., Pruss, S.B., Knoll, A.H., 2009. Evaluating the relationship between the carbon and sulfur cycles in the later Cambrian Ocean: an example from the Port au Port Group, western Newfoundland, Canada. *Earth Planet. Sci. Lett.* 281, 288–297. <http://dx.doi.org/10.1016/j.epsl.2009.02.033>.
- Imhoff, J.F., Trüper, H.G., 1976. Marine sponges as habitats of anaerobic phototrophic bacteria. *Microb. Ecol.* 3, 1–9. <http://dx.doi.org/10.1007/BF02011449>.
- Irwin, M.L., 1965. *General theory of epeiric clear water sedimentation.* AAPG Bull. 49, 445–459.
- James, N.P., Kobluk, D.R., 1978. Lower Cambrian patch reefs and associated sediments: southern Labrador, Canada. *Sedimentology* 25, 1–35. <http://dx.doi.org/10.1111/j.1365-3091.1978.tb00299.x>.
- James, N.P., Wood, R., 2010. Reefs. In: James, N.P., Dalrymple, R.W. (Eds.), *Facies Models 4.* Geological Association of Canada, St. John's, pp. 421–447.
- Jell, J.S., 1984. Cambrian cnidarians with mineralized skeletons. *Palaeontogr. Am.* 54, 105–109.
- Jenkyns, H.C., 1991. Impact of Cretaceous sea level rise and anoxic events on the Mesozoic carbonate platform of Yugoslavia. *AAPG Bull.* 75, 1007–1017. <http://dx.doi.org/10.1306/C9b28ad-1710-11d7-8645000102c1865d>.
- Jenkyns, H.C., 2010. Geochemistry of oceanic anoxic events. *Geochem. Geophys. Geosyst.* 11, Q03004. <http://dx.doi.org/10.1029/2009gc002788>.
- Jenkyns, H.C., Schouten-Huibers, L., Schouten, S., Damsté, J.S.S., 2012. Warm Middle Jurassic-Early Cretaceous high-latitude sea-surface temperatures from the southern ocean. *Clim. Past* 8, 215–226. <http://dx.doi.org/10.5194/cp-8-215-2012>.
- Jiang, S.Y., Pi, D.H., Heubeck, C., Frimmel, H., Liu, Y.P., Deng, H.L., Ling, H.F., Yang, J.H., 2009. Early Cambrian ocean anoxia in South China. *Nature* 459 E5–6; discussion E6. <https://doi.org/10.1038/nature08048>.
- Jin, C., Li, C., Algeo, T.J., Planavsky, N.J., Cui, H., Yang, X., Zhao, Y., Zhang, X., Xie, S., 2016. A highly redox-heterogeneous ocean in South China during the early Cambrian (~529–514 Ma): implications for biota-environment co-evolution. *Earth Planet. Sci. Lett.* 441, 38–51. <http://dx.doi.org/10.1016/j.epsl.2016.02.019>.
- Johns, R.A., Dattilo, B.F., Spincer, B., 2007. Neotype and redescription of the Upper Cambrian anthaspidellid sponge, *Wilbernicyathus donegani* Wilson, 1950. *J. Paleontol.* 81, 435–444. <http://dx.doi.org/10.1666/05028.1>.
- Jourdan, F., Hodges, K., Sell, B., Schaltegger, U., Wingate, M.T.D., Evins, L.Z., Soderlund, U., Haines, P.W., Phillips, D., Blenkinsop, T., 2014. High-precision dating of the Kalkarindji large igneous province, Australia, and synchrony with the Early-Middle Cambrian (Stage 4–5) extinction. *Geology* 42, 543–546. <http://dx.doi.org/10.1130/g35434.1>.
- Kah, L.C., Bartley, J.K., 2011. Protracted oxygenation of the Proterozoic biosphere. *Int. Geol. Rev.* 53, 1424–1442. <http://dx.doi.org/10.1080/00206814.2010.527651>.
- Kah, L.C., Thompson, C.K., Henderson, M.A., Zhan, R., 2016. Behavior of marine sulfur in the Ordovician. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 458, 133–153. <http://dx.doi.org/10.1016/j.palaeo.2015.12.028>.
- Kapp, U.S., 1975. Paleogeology of Middle Ordovician stromatoporeoid mounds in Vermont. *Lethaia* 8, 195–207. <http://dx.doi.org/10.1111/j.1502-3931.1975.tb00923.x>.
- Kasting, J.F., Howard, M.T., Wallmann, K., Veizer, J., Shields, G., Jaffrés, J., 2006. Paleoclimates, ocean depth, and the oxygen isotopic composition of seawater. *Earth Planet. Sci. Lett.* 252, 82–93. <http://dx.doi.org/10.1016/j.epsl.2006.09.029>.
- Keeling, R.E., Körtzinger, A., Gruber, N., 2010. Ocean deoxygenation in a warming world. *Annu. Rev. Mar. Sci.* 2, 199–229. <http://dx.doi.org/10.1146/annurev.marine.010908.163855>.
- Keller, M., Flügel, E., 1996. Early Ordovician reefs from Argentina: Stromatoporeoid vs. Stromatolite origin. *Facies* 34, 177–192. <http://dx.doi.org/10.1007/BF02546163>.
- Kelly, M., 2007. *Porifera: Lithistid Demospongiae (Rock Sponges).* In: NIWA Biodiversity Memoir 121. Graphic Press & Packaging Ltd., Levin (100 pp).
- Kendall, B., Komiya, T., Lyons, T.W., Bates, S.M., Gordon, G.W., Romaniello, S.J., Jiang, G., Creaser, R.A., Xiao, S., McFadden, K., Sawaki, Y., Tahata, M., Shu, D., Han, J., Li, Y., Chu, X., Anbar, A.D., 2015. Uranium and molybdenum isotope evidence for an episode of widespread ocean oxygenation during the late Ediacaran period. *Geochim. Cosmochim. Acta* 156, 173–193. <http://dx.doi.org/10.1016/j.gca.2015.02.025>.
- Kershaw, S., 2015. Paleogeology of the Paleozoic Stromatoporeoida. In: Stearn, C.W. (Ed.), *Treatise on Invertebrate Paleontology, Part E, Porifera, Revised, Hypercalcified Porifera, Volumes 4 and 5.* The University of Kansas Paleontological Institute, Lawrence, Kansas, pp. 631–651.
- Kesling, R.V., Graham, A., 1962. *Ischadites* is a Dasycladacean alga. *J. Paleontol.* 36, 943–952.
- Kiessling, W., 2009. Geologic and biologic controls on the evolution of reefs. *Annu. Rev. Ecol. Evol. Syst.* 40, 173–192. <http://dx.doi.org/10.1146/annurev.ecolsys.110308.120251>.
- Klappa, C.F., James, N.P., 1980. Small lithistid sponge bioherms, early Middle Ordovician Table Head Group, western Newfoundland. *Bull. Can. Petrol. Geol.* 28, 425–451.
- Knight, I., Azmy, K., Boyce, W.D., Lavoie, D., 2008. Tremadocian carbonate rocks of the lower St. George Group, Port au Port Peninsula, western Newfoundland: lithostratigraphic setting of diagenetic, isotopic and geochemistry studies. In: *Geological Survey Report 08–1.* Newfoundland and Labrador Department of Natural Resources, pp. 115–149.
- Knoll, A.H., 2003. Biomineralization and evolutionary history. *Rev. Mineral. Geochem.* 54, 329–356. <http://dx.doi.org/10.2113/0540329>.
- Knoll, A.H., 2011. The multiple origins of complex multicellularity. *Annu. Rev. Earth Planet. Sci.* 39, 217–239. <http://dx.doi.org/10.1146/annurev.earth.031208.100209>.
- Knoll, A.H., Carroll, S.B., 1999. Early animal evolution: emerging views from comparative biology and geology. *Science* 284, 2129–2137. <http://dx.doi.org/10.1126/science.284.5423.2129>.
- Knoll, A.H., Fischer, W.W., 2011. Skeletons and ocean chemistry: the long view. In: Gattuso, J.-P., Hansson, L. (Eds.), *Ocean Acidification.* Oxford University Press, Oxford, pp. 67–82.
- Knoll, A.H., Bambach, R.K., Payne, J.L., Pruss, S., Fischer, W.W., 2007. Paleophysiology and end-Permian mass extinction. *Earth Planet. Sci. Lett.* 256, 295–313. <http://dx.doi.org/10.1016/j.epsl.2007.02.018>.
- Kobluk, D.R., 1979. A new and unusual skeletal organism from the Lower Cambrian of Labrador. *Can. J. Earth Sci.* 16, 2040–2045. <http://dx.doi.org/10.1139/c79-189>.
- Kobluk, D.R., 1984. A new compound skeletal organism from the Rosella Formation (Lower Cambrian), Atan Group, Cassiar Mountains, British Columbia. *J. Paleontol.*

- 58, 703–708.
- Kröger, B., Hints, L., Lehnert, O., 2014. Age, facies, and geometry of the Sandbian/Katian (Upper Ordovician) pelmatozoan-bryozoan-receptaculitid reefs of the Vasalemma Formation, northern Estonia. *Facies* 60, 963–986. <http://dx.doi.org/10.1007/s10347-014-0104-8>.
- Kröger, B., Desrochers, A., Ernst, A., 2017. The reengineering of reef habitats during the Great Ordovician Biodiversification Event. *PALAIOS* 32, 584–599. <http://dx.doi.org/10.2110/palo.2017.017>.
- Kruse, P.D., 1983. Middle Cambrian 'Archaeocyathus' from the Georgina Basin is an anthaspidellid sponge. *Alcheringa* 7, 49–58. <http://dx.doi.org/10.1080/03115518308619633>.
- Kruse, P.D., 1996. Update on the northern Australian Cambrian sponges *Rankenella*, *Jawonya* and *Wagima*. *Alcheringa* 20, 161–178. <http://dx.doi.org/10.1080/03115519608619188>.
- Kruse, P.D., Reitner, J.R., 2014. Northern Australian microbial-metazoan reefs after the mid-Cambrian mass extinction. *Memoirs of the Association of Australasian Palaeontologists* 45, 31–53.
- Kruse, P.D., Zhuravlev, A.Y., 2008. Middle-Late Cambrian *Rankenella-Girvanella* reefs of the Mila Formation, northern Iran. *Can. J. Earth Sci.* 45, 619–639. <http://dx.doi.org/10.1139/e08-016>.
- Kruse, P.D., Zhuravlev, A.Y., James, N.P., 1995. Primordial metazoan-calcimicrobial reefs; Tommotian (Early Cambrian) of the Siberian platform. *PALAIOS* 10, 291–321. <http://dx.doi.org/10.2307/3515157>.
- Kruse, P.D., Gandin, A., Debrénne, F., Wood, R., 1996. Early Cambrian bioconstructions in the Zavkhan Basin of western Mongolia. *Geol. Mag.* 133, 429–444. <http://dx.doi.org/10.1017/S0016756800007597>.
- Kruse, P.D., Zhuravlev, A.Y., Debrénne, F., 2015. Radiocyaths and potentially allied taxa. In: Stearn, C.W. (Ed.), *Treatise on Invertebrate Paleontology, Part E, Porifera, Revised, Hypercalified Porifera, Volumes 4 and 5*. The University of Kansas Paleontological Institute, Lawrence, Kansas, pp. 1085–1093.
- Lafuste, J., Debrénne, F., Gandin, A., Gravestock, D., 1991. The oldest tabulate coral and the associated Archaeocyatha, Lower Cambrian, Flinders Ranges, South Australia. *Geobios* 24, 697–718. [http://dx.doi.org/10.1016/S0016-6995\(06\)80298-6](http://dx.doi.org/10.1016/S0016-6995(06)80298-6).
- Landing, E., 1993. In situ earliest Cambrian tube worms and the oldest metazoan-constructed biostrome (Placentian Series, southeastern Newfoundland). *J. Paleontol.* 67, 333–342. <http://dx.doi.org/10.1017/s0022336000036817>.
- Landing, E., 2012a. The Great American Carbonate Bank in eastern Laurentia: its births, deaths, and linkage to paleoceanic oxygenation (Early Cambrian – Late Ordovician). In: Derby, J.R., Fritz, R.D., Longacre, S.A., Morgan, W.A., Sternbach, C.A. (Eds.), *The Great American Carbonate Bank: The Geology and Economic Resources of the Cambrian-Ordovician Sauk Megasequence of Laurentia*. AAPG Memoir 98. AAPG, pp. 451–492. <http://dx.doi.org/10.1306/13331502M983502>.
- Landing, E., 2012b. Time-specific black mudstones and global hyperwarming on the Cambrian-Ordovician slope and shelf of the Laurentia palaeocontinent. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 367–368, 256–272. <http://dx.doi.org/10.1016/j.palaeo.2011.09.005>.
- Landing, E., English, A., Keppie, J.D., 2010. Cambrian origin of all skeletalized metazoan phyla—discovery of Earth's oldest bryozoans (Upper Cambrian, southern Mexico). *Geology* 38, 547–550. <http://dx.doi.org/10.1130/g30870.1>.
- Landing, E., Antcliffe, J.B., Brasier, M.D., English, A.B., 2015. Distinguishing Earth's oldest known bryozoan (*Pywackia*, late Cambrian) from pennatulacean octocorals (Mesozoic–Recent). *J. Paleontol.* 89, 292–317. <http://dx.doi.org/10.1017/jpa.2014.26>.
- Laub, R.S., 1984. *Lichenaria* Winchell & Schuchert, 1895, *Lamottia* Raymond, 1924 and the early history of the tabulate corals. In: Oliver, W.A., Sando, W.J., Cairns, S.D., Coates, A.G., Macintyre, I.G., Bayer, F.M., Sorauf, J.E. (Eds.), *Recent Advances in the Paleobiology and Geology of the Cnidaria*. *Palaeontographica Americana* Vol. 54. pp. 159–163.
- Lavoie, D., Desrochers, A., Dix, G., Knight, I., Hersi, O.S., 2012. The Great American Carbonate Bank in Eastern Canada: an overview. In: Derby, J.R., Fritz, R.D., Longacre, S.A., Morgan, W.A., Sternbach, C.A. (Eds.), *The Great American Carbonate Bank: The Geology and Economic Resources of the Cambrian-Ordovician Sauk Megasequence of Laurentia*. AAPG memoir 98. AAPG. <http://dx.doi.org/10.1306/13331504M983503>.
- Lavy, A., Keren, R., Yahel, G., Ilan, M., 2016. Intermittent hypoxia and prolonged suboxia measured *in situ* in a marine sponge. *Front. Mar. Sci.* 3, 263. <http://dx.doi.org/10.3389/fmars.2016.00263>.
- Lee, J.-H., Riding, R., 2016. Marine hypoxia, lithistid sponge-microbial communities, and the cause of the Cambrian 'reef gap'. In: 26th Goldschmidt Conference, 1699.
- Lee, J.-H., Chen, J., Choh, S.-J., Lee, D.-J., Han, Z., Chough, S.K., 2014a. Furongian (late Cambrian) sponge-microbial maze-like reefs in the North China platform. *PALAIOS* 29, 27–37. <http://dx.doi.org/10.2110/palo.2013.050>.
- Lee, J.-H., Lee, H.S., Chen, J., Woo, J., Chough, S.K., 2014b. Calcified microbial reefs in the Cambrian Series 2 of the North China Platform: implications for the evolution of Cambrian calcified microbes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 403, 30–42. <http://dx.doi.org/10.1016/j.palaeo.2014.03.020>.
- Lee, M., Sun, N., Choh, S.-J., Lee, D.-J., 2014c. A new Middle Ordovician reef assemblage from north-central China and its palaeobiogeographical implications. *Sediment. Geol.* 310, 30–40. <http://dx.doi.org/10.1016/j.sedgeo.2014.05.003>.
- Lee, J.-H., Chen, J., Chough, S.K., 2015a. The middle-late Cambrian reef transition and related geological events: a review and new view. *Earth Sci. Rev.* 145, 66–84. <http://dx.doi.org/10.1016/j.earscirev.2015.03.002>.
- Lee, J.-H., Chen, J., Woo, J., 2015b. The earliest Phanerozoic carbonate hardground (Cambrian Stage 5, Series 3): implications to the paleoseawater chemistry and early adaptation of hardground fauna. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 440, 172–179. <http://dx.doi.org/10.1016/j.palaeo.2015.07.043>.
- Lee, J.-H., Hong, J., Choh, S.-J., Lee, D.-J., Woo, J., Riding, R., 2016a. Early recovery of sponge framework reefs after Cambrian archaeocyath extinction: Zhangxia Formation (early Cambrian Series 3), Shandong, North China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 457, 269–276. <http://dx.doi.org/10.1016/j.palaeo.2016.06.018>.
- Lee, J.-H., Hong, J., Lee, D.-J., Choh, S.-J., 2016b. A new Middle Ordovician bivalve–siliceous sponge–microbe reef-building consortium from North China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 457, 23–30. <http://dx.doi.org/10.1016/j.palaeo.2016.05.034>.
- Lee, J.-H., Kim, B.-J., Liang, K., Park, T.-Y., Choh, S.-J., Lee, D.-J., Woo, J., 2016c. Cambrian reefs in the western North China platform, Wuhai, Inner Mongolia. *Acta Geol. Sin.* 90, 1946–1954. <http://dx.doi.org/10.1111/1755-6724.13014>.
- Lee, J.-H., Woo, J., Lee, D.-J., 2016d. The earliest reef-building anthaspidellid sponge *Rankenella zhangxianensis* n. sp. from the Zhangxia Formation (Cambrian Series 3), Shandong Province, China. *J. Paleontol.* 90, 1–9. <http://dx.doi.org/10.1017/jpa.2015.53>.
- Leggett, J.K., 1980. British Lower Palaeozoic black shales and their palaeo-oceanographic significance. *J. Geol. Soc. Lond.* 137, 139–156. <http://dx.doi.org/10.1144/gsjgs.137.2.0139>.
- Leinfelder, R.R., 2001. Jurassic reef ecosystems. In: Stanley, G.D. (Ed.), *The History and Sedimentology of Ancient Reef Systems*. Kluwer Academic/Plenum Publishers, New York, pp. 251–309. http://dx.doi.org/10.1007/978-1-4615-1219-6_8.
- Leinfelder, R.R., Krautter, M., Laternser, D.-G.R., Nose, M., Schmid, D.U., Schweigert, G., Werner, W., Keupp, H., Brugger, D.-G.H., Herrmann, R., Rehfeld-Kiefer, U., Schroeder, J.H., Reinhold, C., Koch, R., Zeiss, A., Schweizer, V., Christmann, H., Menges, G., Luterbacher, H., 1994. The origin of Jurassic reefs: current research developments and results. *Facies* 31, 1–56. <http://dx.doi.org/10.1007/BF02536932>.
- Leinfelder, R.R., Werner, W., Nose, M., Schmid, D.U., Krautter, M., Laternser, R., Takacs, M., Hartmann, D., 1996. Paleogeology, growth parameters and dynamics of coral, sponge and microbialite reefs from the Late Jurassic. In: Reitner, J., Neuweiler, F., Gunkel, F. (Eds.), *Global and Regional Controls on Biogenic Sedimentation. 1. Reef Evolution. Research Reports. Göttinger Arbeiten zur Geologie und Paläontologie, Sonderband 2*, Göttingen, pp. 227–248.
- Leinfelder, R.R., Schmid, D.U., Nose, M., Werner, W., 2002. Jurassic reef patterns—the expression of a changing globe. In: Kiessling, W., Flügel, E., Golonka, J. (Eds.), *Phanerozoic Reef Patterns*. SEPM, Tulsa, pp. 465–520. SEPM special publication 72. <https://doi.org/10.2110/pec.02.72.0465>.
- Lenton, T.M., Boyle, R.A., Poulton, S.W., Shields-Zhou, G.A., Butterfield, N.J., 2014. Co-evolution of eukaryotes and ocean oxygenation in the Neoproterozoic era. *Nat. Geosci.* 7, 257–265. <http://dx.doi.org/10.1038/ngeo2108>.
- Lévi, C., 1991. Lithistid sponges from the Norfolk Rise. Recent and Mesozoic genera. In: Reitner, J., Keupp, H. (Eds.), *Fossil and Recent Sponges*. Springer, Berlin, pp. 72–82. http://dx.doi.org/10.1007/978-3-642-75656-6_7.
- Levin, L.A., Ekau, W., Gooday, A.J., Jorissen, F., Middelburg, J.J., Naqvi, S.W.A., Neira, C., Rabalais, N.N., Zhang, J., 2009. Effects of natural and human-induced hypoxia on coastal benthos. *Biogeosciences* 6, 2063–2098. <http://dx.doi.org/10.5194/bg-6-2063-2009>.
- Li, C.-W., Chen, J.-Y., Hua, T.-E., 1998. Precambrian sponges with cellular structures. *Science* 279, 879–882. <http://dx.doi.org/10.1126/science.279.5352.879>.
- Li, Q., Li, Y., Kiessling, W., 2014. Early Ordovician sponge-*Calathium*-microbial reefs on the Yangtze Platform margin of the South China Block. *GFF* 136, 157–161. <http://dx.doi.org/10.1080/11035897.2013.862852>.
- Li, Q., Li, Y., Wang, J., Kiessling, W., 2015. Early Ordovician lithistid sponge-*Calathium* reefs on the Yangtze Platform and their paleoceanographic implications. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 425, 84–96. <http://dx.doi.org/10.1016/j.palaeo.2015.02.034>.
- Li, C., Jin, C., Planavsky, N.J., Algeo, T.J., Cheng, M., Yang, X., Zhao, Y., Xie, S., 2017a. Coupled oceanic oxygenation and metazoan diversification during the early-middle Cambrian? *Geology* 45, 743–746. <http://dx.doi.org/10.1130/g39208.1>.
- Li, Q., Li, Y., Kiessling, W., 2017b. The oldest labechiid stromatolites from in-traskeletal crypts in lithistid sponge-*Calathium* reefs. *Lethaia* 50, 140–148. <http://dx.doi.org/10.1111/let.12182>.
- Li, Q., Li, Y., Zhang, Y., Munnecke, A., 2017c. Dissecting *Calathium*-microbial frameworks: the significance of calathids for the Middle Ordovician reefs in the Tarim Basin, northwestern China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 474, 66–78. <http://dx.doi.org/10.1016/j.palaeo.2016.08.005>.
- Liu, B., Rigby, J.K., Zhu, Z., 2003. Middle Ordovician lithistid sponges from the Bachukalpina area, Xinjiang, northwestern China. *J. Paleontol.* 77, 430–441. [http://dx.doi.org/10.1666/0022-3360\(2003\)077<0430:molsft>2.0.co;2](http://dx.doi.org/10.1666/0022-3360(2003)077<0430:molsft>2.0.co;2).
- Loch, J.D., Stitt, J.H., Derby, J.R., 1993. Cambrian-Ordovician boundary interval extinctions: implications of revised trilobite and brachiopod data from Mount Wilson, Alberta, Canada. *J. Paleontol.* 67, 497–517. <http://dx.doi.org/10.1017/S0022336000024859>.
- Love, G.D., Grosjean, E., Stalvies, C., Fike, D.A., Grotzinger, J.P., Bradley, A.S., Kelly, A.E., Bhatia, M., Meredith, W., Snape, C.E., Bowring, S.A., Condon, D.J., Summons, R.E., 2009. Fossil steroids record the appearance of Demospongiae during the Cryogenian period. *Nature* 457, 718–721. <http://dx.doi.org/10.1038/nature07673>.
- Luo, C., Reitner, J., 2014. First report of fossil "keratose" demosponges in Phanerozoic carbonates: preservation and 3-D reconstruction. *Naturwissenschaften* 101, 467–477. <http://dx.doi.org/10.1007/s00114-014-1176-0>.
- Ma, J., Taylor, P.D., Xia, F., Zhan, R., Sevastopulo, G., 2015. The oldest known bryozoan: *Prophyllodictya* (Cryptostomata) from the lower Tremadocian (Lower Ordovician) of Liujiaochang, south-western Hubei, central China. *Palaeontology* 58, 925–934. <http://dx.doi.org/10.1111/pala.12189>.
- Maldonado, M., Aguilar, R., Blanco, J., Garcia, S., Serrano, A., Punzon, A., 2015. Aggregated clumps of lithistid sponges: a singular, reef-like bathyal habitat with relevant paleontological connections. *PLoS One* 10, e0125378. <http://dx.doi.org/10.1371/journal.pone.0125378>.
- Maldonado, M., Aguilar, R., Bannister, R.J., Bell, J.J., Conway, K.W., Dayton, P.K., Díaz, C., Gutt, J., Kelly, M., Kenchington, E.L.R., Leys, S.P., Pomponi, S.A., Rapp, H.T., Rützler, K., Tendal, O.S., Vacelet, J., Young, C.M., 2016. Sponge grounds as key marine habitats: a synthetic review of types, structure, functional roles, and conservation concerns. In: Rossi, S., Bramanti, L., Gori, A., Orejas, C. (Eds.), *Marine Animal Forests: The Ecology of Benthic Biodiversity Hotspots*. Springer, Cham, pp.

- 1–39. http://dx.doi.org/10.1007/978-3-319-17001-5_24-1.
- Maloof, A.C., Porter, S.M., Moore, J.L., Dudas, F.O., Bowring, S.A., Higgins, J.A., Fike, D.A., Eddy, M.P., 2010a. The earliest Cambrian record of animals and ocean geochemical change. *Geol. Soc. Am. Bull.* 122, 1731–1774. <http://dx.doi.org/10.1130/b30346.1>.
- Maloof, A.C., Rose, C.V., Beach, R., Samuels, B.M., Calmet, C.C., Erwin, D.H., Poirier, G.R., Yao, N., Simons, F.J., 2010b. Possible animal-body fossils in pre-Marinoan limestones from South Australia. *Nat. Geosci.* 3, 653–659. <http://dx.doi.org/10.1038/ngeo934>.
- Marenco, P.J., Martin, K.R., Marenco, K.N., Barber, D.C., 2016. Increasing global ocean oxygenation and the Ordovician radiation: insights from Th/U of carbonates from the Ordovician of western Utah. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 458, 77–84. <http://dx.doi.org/10.1016/j.palaeo.2016.05.014>.
- Markello, J.R., Read, J.F., 1981. Carbonate ramp-to-deeper shale shelf transitions of an Upper Cambrian intrashelf basin, Nolichucky Formation, Southwest Virginia Appalachians. *Sedimentology* 28, 573–597. <http://dx.doi.org/10.1111/j.1365-3091.1981.tb01702.x>.
- Matabos, M., Tunnicliffe, V., Juniper, S.K., Dean, C., 2012. A year in hypoxia: epibenthic community responses to severe oxygen deficit at a subsea observatory in a coastal inlet. *PLoS One* 7, e45626. <http://dx.doi.org/10.1371/journal.pone.0045626>.
- McKenzie, N.R., Hughes, N.C., Gill, B.C., Myrow, P.M., 2014. Plate tectonic influences on Neoproterozoic-early Paleozoic climate and animal evolution. *Geology* 42, 127–130. <http://dx.doi.org/10.1130/g34962.1>.
- Mentel, M., Rottger, M., Leys, S., Tielens, A.G., Martin, W.F., 2014. Of early animals, anaerobic mitochondria, and a modern sponge. *BioEssays* 36, 924–932. <http://dx.doi.org/10.1002/bies.201400060>.
- Mi, J., Zhang, S., Chen, J., Tang, L., He, Z., 2007. The distribution of the oil derived from Cambrian source rocks in Lunnan area, the Tarim Basin, China. *Chin. Sci. Bull.* 52, 133–140. <http://dx.doi.org/10.1007/s11434-007-6004-x>.
- Middleburg, J.J., Levin, L.A., 2009. Coastal hypoxia and sediment biogeochemistry. *Biogeochemistry* 6, 1273–1293. <http://dx.doi.org/10.5194/bg-6-1273-2009>.
- Miller, K.G., Kominz, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E., Sugarman, P.J., Cramer, B.S., Christie-Blick, N., Pekar, S.F., 2005. The Phanerozoic record of global sea-level change. *Science* 310, 1293–1298. <http://dx.doi.org/10.1126/science.1116412>.
- Miller, J.F., Evans, K.R., Dattilo, B.F., 2012. The Great American Carbonate Bank in the miogeoclinal of western central Utah: tectonic influences on sedimentation. In: Derby, J.R., Fritz, R.D., Longacre, S.A., Morgan, W.A., Sternbach, C.A. (Eds.), *The Great American Carbonate Bank: The Geology and Economic Resources of the Cambrian-Ordovician Sauk Megasequence of Laurentia*. AAPG Memoir 98. AAPG, pp. 769–854. <http://dx.doi.org/10.1306/13331516M983498>.
- Mills, D.B., Canfield, D.E., 2014. Oxygen and animal evolution: did a rise of atmospheric oxygen “trigger” the origin of animals? *BioEssays* 36, 1145–1155. <http://dx.doi.org/10.1002/bies.201400101>.
- Mills, D.B., Ward, L.M., Jones, C., Sweeten, B., Forth, M., Treusch, A.H., Canfield, D.E., 2014. Oxygen requirements of the earliest animals. *Proc. Natl. Acad. Sci.* 111, 4168–4172. <http://dx.doi.org/10.1073/pnas.1400547111>.
- Montañez, I.P., Osleger, D.A., Banner, J.L., Mack, L.E., Musgrove, M., 2000. Evolution of the Sr and C isotope composition of Cambrian oceans. *GSA Today* 10, 1–5.
- Morgan, W.A., 2012. Sequence stratigraphy of the Great American Carbonate Bank. In: Derby, J.R., Fritz, R.D., Longacre, S.A., Morgan, W.A., Sternbach, C.A. (Eds.), *The Great American Carbonate Bank: The Geology and Economic Resources of the Cambrian-Ordovician Sauk Megasequence of Laurentia*. AAPG Memoir 98. AAPG, pp. 37–82. <http://dx.doi.org/10.1306/13331499M980271>.
- Morrow, C., Cárdenas, P., 2015. Proposal for a revised classification of the Demospongiae (Porifera). *Front. Zool.* 12, 7. <http://dx.doi.org/10.1186/s12983-015-0099-8>.
- Mrozek, S., Dattilo, B.F., Hicks, M., Miller, J.F., 2003. Metazoan reefs from the Upper Cambrian of the Arrow Canyon Range, Clark County, Nevada. *Geol. Soc. Am. Abstr. Programs* 35, 500.
- Mundy, D.J.C., 1994. Microbialite-sponge-bryozoan-coral framestones in Lower Carboniferous (Late Viséan) buildups of northern England (UK). In: Embry, A.F., Beauchamp, B., Glass, D.J. (Eds.), *Pangea: Global Environments and Resources*. CSPG Special Publications 17. Canadian Society of Petroleum Geologists, Calgary, pp. 713–729.
- Muscente, A.D., Marc Michel, F., Dale, J.G., Xiao, S., 2015. Assessing the veracity of Precambrian ‘sponge’ fossils using *in situ* nanoscale analytical techniques. *Precambrian Res.* 263, 142–156. <http://dx.doi.org/10.1016/j.precamres.2015.03.010>.
- Myagkova, Y.I., 1966. Soanites—a new group of organisms. *Int. Geol. Rev.* 8, 795–802. <http://dx.doi.org/10.1080/00206816609474340>.
- Na, L., Kiessling, W., 2015. Diversity partitioning during the Cambrian radiation. *Proc. Natl. Acad. Sci.* 112, 4702–4706. <http://dx.doi.org/10.1073/pnas.1424985112>.
- Nance, R.D., Murphy, J.B., 2013. Origins of the supercontinent cycle. *Geosci. Front.* 4, 439–448. <http://dx.doi.org/10.1016/j.gsf.2012.12.007>.
- Newell, N.D., 1972. The evolution of reefs. *Sci. Am.* 226, 54–69.
- Nitecki, M.H., 1970. North American cyclocrinid algae. In: *Fieldiana Geology* 21. History, Field Museum of Natural (182 pp).
- Nitecki, M.H., 1972. *Ischadites abbottiae*, a new North American Silurian species (Dasycladales). *Phycologia* 10, 263–275.
- Nitecki, M.H., Debrenne, F., 1979. The nature of Radiocyathids and their relationship to Receptaculitids and Archaeocyathids. *Geobios* 12, 5–27. [http://dx.doi.org/10.1016/S0016-6995\(79\)80106-0](http://dx.doi.org/10.1016/S0016-6995(79)80106-0).
- Nitecki, M.H., Mutvei, H., Nitecki, D.V., 1999. Receptaculitids: a phylogenetic debate on a problematic fossil taxon. *Kluwer/Plenum, New York* (241 pp).
- Nitecki, M.H., Webby, B.D., Spjelldnaes, N., Zhen, Y.-Y., 2004. Receptaculitids and algae. In: Webby, B.D., Paris, F., Droser, M.L., Percival, I.G. (Eds.), *The Great Ordovician Biodiversification Event*. Columbia University Press, New York, pp. 336–347.
- Nowak, H., Servais, T., Monnet, C., Molyneux, S.G., Vandenbroucke, T.R.A., 2015. Phytoplankton dynamics from the Cambrian Explosion to the onset of the Great Ordovician Biodiversification Event: a review of Cambrian acritarch diversity. *Earth Sci. Rev.* 151, 117–131. <http://dx.doi.org/10.1016/j.earscirev.2015.09.005>.
- Och, L.M., Cremonese, L., Shields-Zhou, G.A., Poulton, S.W., Struck, U., Ling, H., Li, D., Chen, X., Manning, C., Thirlwall, M., Strauss, H., Zhu, M., Fairchild, I., 2016. Palaeoceanographic controls on spatial redox distribution over the Yangtze platform during the Ediacaran-Cambrian transition. *Sedimentology* 63, 378–410. <http://dx.doi.org/10.1111/sed.12220>.
- Ogg, J.G., Ogg, G., Gradstein, F.M., 2016. A concise geologic time scale. *Elsevier* (240 pp).
- Oliver, W.A., 1996. Origins and relationships of Paleozoic coral groups and the origin of the Scleractinia. *Paleontological Society Papers* 1, 107–134.
- Oliver, W.A., Coates, A.G., 1987. *Phylum Cnidaria*. In: Boardman, R.S., Cheetham, A.H., Rowell, A.J. (Eds.), *Fossil Invertebrates*. Blackwell, Oxford, pp. 140–193.
- Opalinski, P.R., Harland, T.L., 1981. The Middle Ordovician of the Oslo Region, Norway, 29. Stratigraphy of the Mjøsa Limestone in the Toten and Nes-Hamar areas. *Nor. Geol. Tidsskr.* 61, 59–78.
- Pacheco, M.L., Galante, D., Rodrigues, F., Leme Jde, M., Bidola, P., Hagadorn, W., Stockmar, M., Herzen, J., Rudnitzki, I.D., Pfeiffer, F., Marques, A.C., 2015. Insights into the skeletonization, lifestyle, and affinity of the unusual Ediacaran fossil *Corumbella*. *PLoS One* 10, e0114219. <http://dx.doi.org/10.1371/journal.pone.0114219>.
- Pages, A., Schmid, S., 2016. Euxinia linked to the Cambrian Drumian carbon isotope excursion (DICE) in Australia: geochemical and chemostratigraphic evidence. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 461, 65–76. <http://dx.doi.org/10.1016/j.palaeo.2016.08.008>.
- Palmer, A.R., 1984. The biore problem: evolution of an idea. *J. Paleontol.* 58, 599–611.
- Palmer, A.R., James, N.P., 1979. The Hawke Bay event: A circum-lapetus regression near the lower Middle Cambrian boundary. In: Wones, D.R. (Ed.), *The Caledonides in the U.S.A. I.G.C.P. Project 27: Caledonide orogen*. Department of Geological Sciences, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, pp. 15–18.
- Parente, M., Frijia, G., Di Lucia, M., Jenkyns, H.C., Woodfine, R.G., Baroncini, F., 2008. Stepwise extinction of larger foraminifers at the Cenomanian-Turonian boundary: a shallow-water perspective on nutrient fluctuations during oceanic anoxic event 2 (Bonarelli event). *Geology* 36, 715–718. <http://dx.doi.org/10.1130/g24893a.1>.
- Park, T.-Y., Woo, J., Lee, D.-J., Lee, D.-C., Lee, S.-B., Han, Z., Chough, S.K., Choi, D.K., 2011. A stem-group cnidarian described from the mid-Cambrian of China and its significance for cnidarian evolution. *Nat. Commun.* 2, 442. <http://dx.doi.org/10.1038/ncomms1457>.
- Park, T.-Y.S., Kihm, J.-H., Woo, J., Zhen, Y.-Y., Engelbretsen, M., Hong, J., Choh, S.-J., Lee, D.-J., 2016. Cambrian stem-group cnidarians with a new species from the Cambrian Series 3 of the Taebaeksan Basin, Korea. *Acta Geol. Sin.* 90, 827–837. <http://dx.doi.org/10.1111/1755-6724.12726>.
- Pashin, J.C., Kopaska-Merkel, D.C., Arnold, A.C., McIntyre, M.R., Thomas, W.A., 2012. Gigantic, gaseous mudwads in Cambrian shale: Conasauga Formation, southern Appalachians, USA. *Int. J. Coal Geol.* 103, 70–91. <http://dx.doi.org/10.1016/j.coal.2012.05.010>.
- Pavela, J.S., Ross, J.L., Chittenden, M.E., 1983. Sharp reductions in abundance of fishes and benthic macroinvertebrates in the Gulf of Mexico off Texas associated with hypoxia. *Northeast Gulf Sci.* 6, 167–173.
- Payne, J.L., Boyer, A.G., Brown, J.H., Finnegan, S., Kowalewski, M., Krause Jr., R.A., Lyons, S.K., McClain, C.R., McShea, D.W., Novack-Gottshall, P.M., Smith, F.A., Stempian, J.A., Wang, S.C., 2009. Two-phase increase in the maximum size of life over 3.5 billion years reflects biological innovation and environmental opportunity. *Proc. Natl. Acad. Sci.* 106, 24–27. <http://dx.doi.org/10.1073/pnas.0806314106>.
- Payne, J.L., McClain, C.R., Boyer, A.G., Brown, J.H., Finnegan, S., Kowalewski, M., Krause Jr., R.A., Lyons, S.K., McShea, D.W., Novack-Gottshall, P.M., Smith, F.A., Spaeth, P., Stempian, J.A., Wang, S.C., 2011. The evolutionary consequences of oxygenic photosynthesis: a body size perspective. *Photosynth. Res.* 107, 37–57. <http://dx.doi.org/10.1007/s11120-010-9593-1>.
- Pedersen, T.F., Calvert, S.E., 1990. Anoxia vs. productivity: what controls the formation of organic-carbon-rich sediments and sedimentary rocks? *AAPG Bull.* 74, 454–466.
- Peel, J.S., 2011. The coral *Cothonion* from the lower Cambrian of North Greenland. *Alcheringa* 35, 405–411. <http://dx.doi.org/10.1080/03115518.2011.521438>.
- Peel, J.S., 2017a. The oldest pelmatozoan encrusted hardground and holdfasts from Laurentia (Cambrian Series 2–3). *GFF* 139, 195–204. <http://dx.doi.org/10.1080/11035897.2017.1347196>.
- Peel, J.S., 2017b. A problematic cnidarian (*Cambroctoconus*; Octocorallia?) from the Cambrian (Series 2–3) of Laurentia. *J. Paleontol.* 91, 871–882. <http://dx.doi.org/10.1017/jpa.2017.49>.
- Peng, S., Geyer, G., Hamdi, B., 1999. Trilobites from the Shahmirzad section, Alborz Mountains, Iran: their taxonomy, biostratigraphy and bearing for international correlation. *Beringeria* 25, 3–66.
- Peng, S., Babcock, L.E., Cooper, R.A., 2012. The Cambrian Period. In: Gradstein, F.M., Ogg, J.G., Schmitz, M., Ogg, G. (Eds.), *The Geologic Time Scale 2012*. Elsevier, pp. 437–488. <http://dx.doi.org/10.1016/b978-0-444-59425-9.00019-6>.
- Penny, A.M., Wood, R., Curtis, A., Bowyer, F., Tostevin, R., Hoffman, K.H., 2014. Ediacaran metazoan reefs from the Nama Group, Namibia. *Science* 344, 1504–1506. <http://dx.doi.org/10.1126/science.1253393>.
- Penny, A.M., Wood, R.A., Zhuravlev, A.Y., Curtis, A., Bowyer, F., Tostevin, R., 2017. Intraspecific variation in an Ediacaran skeletal metazoan: *Namacalathus* from the Nama Group, Namibia. *Geobiology* 15, 81–93. <http://dx.doi.org/10.1111/gbi.12205>.
- Percival, L.M.E., Witt, M.L.L., Mather, T.A., Hermoso, M., Jenkyns, H.C., Hesselbo, S.P., Al-Suwaidi, A.H., Storm, M.S., Xu, W., Ruhl, M., 2015. Globally enhanced mercury deposition during the end-Plensbachian extinction and Toarcian OAE: a link to the Karoo–Ferrar large Igneous Province. *Earth Planet. Sci. Lett.* 428, 267–280. <http://dx.doi.org/10.1016/j.epsl.2015.06.064>.
- Perfetta, P.J., Shelton, K.L., Stitt, J.H., 1999. Carbon isotope evidence for deep-water invasion at the Marjumiid-Pteroccephaliid biore boundary, Black Hills, USA: a common origin for biotic crises on Late Cambrian shelves. *Geology* 27, 403. [http://dx.doi.org/10.1130/0091-7613\(1999\)027<0403:ciefdw>2.3.co;2](http://dx.doi.org/10.1130/0091-7613(1999)027<0403:ciefdw>2.3.co;2).
- Peters, S.E., Gaines, R.R., 2012. Formation of the ‘Great Unconformity’ as a trigger for the

- Cambrian explosion. *Nature* 484, 363–366. <http://dx.doi.org/10.1038/nature10969>.
- Pickett, J.W., Percival, I.G., 2001. Ordovician faunas and biostratigraphy in the Gunningbland area, central New South Wales, Australia. *Alcheringa* 25, 9–52. <http://dx.doi.org/10.1080/03115511008619212>.
- Pisera, A., 2002. Fossil 'Lithistids': An Overview. In: Hooper, J.N.A., Van Soest, R.W.M. (Eds.), *Systema Porifera: A Guide to the Classification of Sponges*. Kluwer Academic/Plenum Publishers, New York, pp. 388–402.
- Pisera, A., Lévi, C., 2002. 'Lithistid' Demospongiae. In: Hooper, J.N.A., Van Soest, R.W.M. (Eds.), *Systema Porifera: A Guide to the Classification of Sponges*. Kluwer Academic/Plenum Publishers, New York, pp. 299–301.
- Pitcher, M., 1964. Evolution of Chazy (Ordovician) reefs of eastern United States and Canada. *Bull. Can. Petrol. Geol.* 12, 632–691.
- Pool, W., Geluk, M., Abels, J., Tiley, G., Idiz, E., Leenaerts, E., 2012. Assessment of an unusual European shale gas play: the Cambro-Ordovician Alum Shale, southern Sweden. *Proceedings of the Society of Petroleum Engineers/European Association of Geoscientists and Engineers Unconventional Resources Conference*, Vienna, pp. 152339.
- Praetorius, S.K., Mix, A.C., Walczak, M.H., Wolhowe, M.D., Addison, J.A., Prah, F.G., 2015. North Pacific deglacial hypoxic events linked to abrupt ocean warming. *Nature* 527, 362–366. <http://dx.doi.org/10.1038/nature15753>.
- Pratt, B.R., 1989. Small early Middle Ordovician patch reefs, Laval Formation (Chazy Group), Caughnawaga, Montreal area, Quebec. In: Geldsetzer, H.H.J., James, N.P., Tebbutt, G.E. (Eds.), *Reefs: Canada and adjacent areas*. Canadian Society of Petroleum Geologists Memoir 13. Canadian Society of Petroleum Geologists, pp. 218–223.
- Pratt, B.R., James, N.P., 1982. Cryptalgal-metazoan bioherms of early Ordovician age in the St George Group, western Newfoundland. *Sedimentology* 29, 543–569. <http://dx.doi.org/10.1111/j.1365-3091.1982.tb01733.x>.
- Pratt, B.R., Spincer, B.R., Wood, R.A., Zhuravlev, A.Y., 2001. Ecology and evolution of Cambrian reefs. In: Zhuravlev, A.Y., Riding, R. (Eds.), *Ecology of the Cambrian Radiation*. Columbia University Press, New York, pp. 254–274.
- Pruss, S.B., Knoll, A.H., 2017. Environmental covariation of metazoans and microbialites in the Lower Ordovician Boat Harbour Formation, Newfoundland. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 485, 917–929. <http://dx.doi.org/10.1016/j.palaeo.2017.08.007>.
- Pruss, S.B., Finnegan, S., Fischer, W.W., Knoll, A.H., 2010. Carbonates in skeleton-poor seas: new insights from Cambrian and Ordovician strata of Laurentia. *PALAIOS* 25, 73–84. <http://dx.doi.org/10.2110/palo.2009.p09-101r>.
- Quinn, N.J., Kojis, B.L., 1999. Community structure of the living fossil coralline sponge populations at the Grotto, Saipan, Northern Mariana Islands. *Bull. Mar. Sci.* 65, 227–234.
- Rabalais, N.N., Díaz, R.J., Levin, L.A., Turner, R.E., Gilbert, D., Zhang, J., 2010. Dynamics and distribution of natural and human-caused hypoxia. *Biogeosciences* 7, 585–619. <http://dx.doi.org/10.5194/bg-7-585-2010>.
- Railsback, L.B., Ackerly, S.C., Anderson, T.F., Cisne, J.L., 1990. Palaeontological and isotope evidence for warm saline deep waters in Ordovician oceans. *Nature* 343, 156–159. <http://dx.doi.org/10.1038/343156a0>.
- Rasmussen, C.M., Ullmann, C.V., Jakobsen, K.G., Lindskog, A., Hansen, J., Hansen, T., Eriksson, M.E., Dronov, A., Frei, R., Korte, C., Nielsen, A.T., Harper, D.A., 2016. Onset of main Phanerozoic marine radiation sparked by emerging mid Ordovician icehouse. *Sci. Rep.* 6, 18884. <http://dx.doi.org/10.1038/srep18884>.
- Reitner, J., 1993. Modern cryptic microbialite/metazoan facies from Lizard Island (Great Barrier Reef, Australia) formation and concepts. *Facies* 29, 3–40. <http://dx.doi.org/10.1007/BF02536915>.
- Renaud, M.L., 1986. Detecting and avoiding oxygen deficient sea water by brown shrimp, *Penaeus aztecus* (Ives), and white shrimp *Penaeus setiferus* (Linnaeus). *J. Exp. Mar. Biol. Ecol.* 98, 283–292. [http://dx.doi.org/10.1016/0022-0981\(86\)90218-2](http://dx.doi.org/10.1016/0022-0981(86)90218-2).
- Rhoads, D.C., Morse, J.W., 1971. Evolutionary and ecologic significance of oxygen-deficient marine basins. *Lethaia* 4, 413–428. <http://dx.doi.org/10.1111/j.1502-3931.1971.tb01864.x>.
- Riding, R., 2001. Calcified algae and bacteria. In: Zhuravlev, A.Y., Riding, R. (Eds.), *Ecology of the Cambrian Radiation*. Columbia University Press, New York, pp. 445–473.
- Riding, R., 2002. Structure and composition of organic reefs and carbonate mud mounds: concepts and categories. *Earth Sci. Rev.* 58, 163–231. [http://dx.doi.org/10.1016/S0012-8252\(01\)00089-7](http://dx.doi.org/10.1016/S0012-8252(01)00089-7).
- Riding, R., 2004. *Solenopora* is a chaetetid sponge, not an alga. *Palaeontology* 47, 117–122. <http://dx.doi.org/10.1111/j.0031-0239.2004.00351.x>.
- Riding, R., 2006. Microbial carbonate abundance compared with fluctuations in metazoan diversity over geological time. *Sediment. Geol.* 185, 229–238. <http://dx.doi.org/10.1016/j.sedgeo.2005.12.015>.
- Riding, R., Voronova, L., 1984. Assemblages of calcareous algae near the Precambrian/Cambrian boundary in Siberia and Mongolia. *Geol. Mag.* 121, 205–210. <http://dx.doi.org/10.1017/S0016756800028260>.
- Riding, R., Zhuravlev, A.Y., 1994. Cambrian reef builders: calcimicrobes and archaeocyaths, ecological aspects of the Cambrian Radiation, IGCP Project 366. *Terra Nova*, Cambridge, pp. 7.
- Riding, R., Zhuravlev, A.Y., 1995. Structure and diversity of oldest sponge-microbe reefs: Lower Cambrian, Aldan River, Siberia. *Geology* 23, 649–652. [http://dx.doi.org/10.1130/0091-7613\(1995\)023<0649:SADOOS>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1995)023<0649:SADOOS>2.3.CO;2).
- Riedel, B., Pados, T., Pretterebner, K., Schiemer, L., Steckbauer, A., Haselmair, A., Zuschin, M., Stachowitsch, M., 2014. Effect of hypoxia and anoxia on invertebrate behaviour: ecological perspectives from species to community level. *Biogeosciences* 11, 1491–1518. <http://dx.doi.org/10.5194/bg-11-1491-2014>.
- Rietschel, L.S., 1977. Receptaculitids are calcareous algae but no dasyclads. In: Flügel, E. (Ed.), *Fossil Algae: Recent Results and Developments*. Springer, Berlin Heidelberg, Berlin, Heidelberg, pp. 212–214. http://dx.doi.org/10.1007/978-3-642-66516-5_22.
- Rietschel, S., Nitecki, M.H., 1984. Ordovician receptaculitid algae from Burma. *Palaeontology* 27, 415–420.
- Rowland, S.M., 1984. Were there framework reefs in the Cambrian? *Geology* 12, 181–183. [http://dx.doi.org/10.1130/0091-7613\(1984\)12<181:wtfrt>2.0.co;2](http://dx.doi.org/10.1130/0091-7613(1984)12<181:wtfrt>2.0.co;2).
- Rowland, S.M., 2001. Archaeocyaths—a history of phylogenetic interpretation. *J. Paleontol.* 75, 1065–1078. [http://dx.doi.org/10.1666/0022-3366\(2001\)075<1065:AAHOPI>2.0.CO;2](http://dx.doi.org/10.1666/0022-3366(2001)075<1065:AAHOPI>2.0.CO;2).
- Rowland, S.M., Gangloff, R.A., 1988. Structure and paleoecology of Lower Cambrian reefs. *PALAIOS* 3, 111–135. <http://dx.doi.org/10.2307/3514525>.
- Rowland, S.M., Shapiro, R.S., 2002. Reef patterns and environmental influences in the Cambrian and earliest Ordovician. In: Kiessling, W., Flügel, E., Golonka, J. (Eds.), *Phanerozoic Reef Patterns*. SEPM, Tulsa, pp. 95–128 SEPM special publication 72.
- Royer, D.L., Berner, R.A., Montañez, I.P., Tabor, N.J., Beerling, D.J., 2004. CO₂ as a primary driver of Phanerozoic climate. *GSA Today* 14, 4–10. [http://dx.doi.org/10.1130/1052-5173\(2004\)014<4:caapdo>2.0.co;2](http://dx.doi.org/10.1130/1052-5173(2004)014<4:caapdo>2.0.co;2).
- Rozanov, A.Y., Zhuravlev, A.Y., 1992. The Lower Cambrian fossil record of the Soviet Union. In: Lipps, J.H., Signor, P.W. (Eds.), *Origin and Early Evolution of the Metazoa*. Plenum Press, New York, pp. 205–282.
- Ryder, R.T., Harris, D.C., Gerome, P., Hainsworth, T.J., Burruss, R.A., Lillis, P.G., Jarvie, D.M., Pawlewicz, M.J., 2014. Evidence for Cambrian petroleum source rocks in the Rome trough of West Virginia and Kentucky, Appalachian basin. In: *Coal and Petroleum Resources in the Appalachian Basin: Distribution, Geologic Framework, and Geochemical Character*. US Geological Survey, pp. 36. <http://dx.doi.org/10.3133/pp1708G.8>.
- Sahoo, S.K., Planavsky, N.J., Kendall, B., Wang, X., Shi, X., Scott, C., Anbar, A.D., Lyons, T.W., Jiang, G., 2012. Ocean oxygenation in the wake of the Marinoan glaciation. *Nature* 489, 546–549. <http://dx.doi.org/10.1038/nature11445>.
- Saltzman, M.R., 1999. Upper Cambrian carbonate platform evolution, *Elvinia* and *Taenicephalus* zones (Pteroccephaliid-Ptychaspis biomere boundary), northwestern Wyoming. *J. Sediment. Res.* 69, 926–938. <http://dx.doi.org/10.2110/jsr.69.926>.
- Saltzman, M.R., 2005. Phosphorus, nitrogen, and the redox evolution of the Paleozoic oceans. *Geology* 33, 573–576. <http://dx.doi.org/10.1130/g21535.1>.
- Saltzman, M.R., Davidson, J.P., Holden, P., Runnegar, B., Lohmann, K.C., 1995. Sea-level-driven changes in ocean chemistry at an Upper Cambrian extinction horizon. *Geology* 23, 893–896. [http://dx.doi.org/10.1130/0091-7613\(1995\)023<0893:SLDCIO>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1995)023<0893:SLDCIO>2.3.CO;2).
- Saltzman, M.R., Ripperdan, R.L., Brasier, M.D., Lohmann, K.C., Robison, R.A., Chang, W.T., Peng, S., Ergaliev, E.K., Runnegar, B., 2000. A global carbon isotope excursion (SPICE) during the Late Cambrian: relation to trilobite extinctions, organic-matter burial and sea level. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 162, 211–223. [http://dx.doi.org/10.1016/S0031-0182\(00\)00128-0](http://dx.doi.org/10.1016/S0031-0182(00)00128-0).
- Saltzman, M.R., Young, S.A., Kump, L.R., Gill, B.C., Lyons, T.W., Runnegar, B., 2011. Pulse of atmospheric oxygen during the late Cambrian. *Proc. Natl. Acad. Sci.* 162, 3876–3881. <http://dx.doi.org/10.1073/pnas.1011836108>.
- Saltzman, M.R., Edwards, C.T., Adrain, J.M., Westrop, S.R., 2015. Persistent oceanic anoxia and elevated extinction rates separate the Cambrian and Ordovician radiations. *Geology* 43, 807–810. <http://dx.doi.org/10.1130/g36814.1>.
- Savarese, M., Mount, J.F., Sorauf, J.E., Bucklin, L., 1993. Paleobiologic and paleoenvironmental context of coral-bearing Early Cambrian reefs: implications for Phanerozoic reef development. *Geology* 21, 917–920. [http://dx.doi.org/10.1130/0091-7613\(1993\)021\(0917:PAPCOC\)2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1993)021(0917:PAPCOC)2.3.CO;2).
- Schlanger, S.O., Jenkyns, H.C., 1976. Cretaceous Oceanic Anoxic Events: causes and consequences. *Geol. Mijnb.* 55, 179–184.
- Schlappé, M.L., Schöttner, S.I., Lavik, G., Kuypers, M.M., de Beer, D., Hoffmann, F., 2010. Evidence of nitrification and denitrification in high and low microbial abundance sponges. *Mar. Biol.* 157, 593–602. <http://dx.doi.org/10.1007/s00227-009-1344-5>.
- Schmitt, S., Tsai, P., Bell, J., Fromont, J., Ilan, M., Lindquist, N., Perez, T., Rodrigo, A., Schupp, P.J., Vacelet, J., Webster, N., Hentschel, U., Taylor, M.W., 2012. Assessing the complex sponge microbiota: core, variable and species-specific bacterial communities in marine sponges. *ISME J.* 6, 564–576. <http://dx.doi.org/10.1038/ismej.2011.116>.
- Schmittner, A., Oeschlies, A., Matthews, H.D., Galbraith, E.D., 2008. Future changes in climate, ocean circulation, ecosystems, and biogeochemical cycling simulated for a business-as-usual CO₂ emission scenario until year 4000 AD. *Glob. Biogeochem. Cycles* 22, GB1013. <http://dx.doi.org/10.1029/2007gb002953>.
- Schovsbo, N.H., Nielsen, A.T., Klitten, K., Mathiesen, A., Rasmussen, P., 2011. Shale gas investigations in Denmark: Lower Palaeozoic shales on Bornholm. *Geol. Surv. Denmark Greenland Bull.* 23, 9–12.
- Schröder, S., Grotzinger, J.P., 2007. Evidence for anoxia at the Ediacaran–Cambrian boundary: the record of redox-sensitive trace elements and rare earth elements in Oman. *J. Geol. Soc.* 164, 175–187. <http://dx.doi.org/10.1144/0016-76492005-022>.
- Schuster, A., Erpenbeck, D., Pisera, A., Hooper, J., Bryce, M., Fromont, J., Wörheide, G., 2015. Deceptive desmas: molecular phylogenetics suggests a new classification and uncovers convergent evolution of lithistid demosponges. *PLoS One* 10, e116038. <http://dx.doi.org/10.1371/journal.pone.0116038>.
- Scrutton, C.T., 1979. Early fossil cnidarians. In: House, M.R. (Ed.), *The Origin of Major Invertebrate Groups*. Academic, London, pp. 161–207.
- Scrutton, C.T., 1992. *Flindersipora bowmani* Lafuste, and the early evolution of the tabulate corals. *Fossil Cnidaria Porifera* 21, 29–33.
- Scrutton, C.T., 1997. The Palaeozoic corals, I: origins and relationships. *Proc. Yorks. Geol. Soc.* 51, 177–208. <http://dx.doi.org/10.1144/pygs.51.3.177>.
- Scrutton, C.T., 1999. Palaeozoic corals: their evolution and palaeoecology. *Geol. Today* 15, 184–193. <http://dx.doi.org/10.1046/j.1365-2451.1999.1505005.x>.
- Seilacher, A., 1999. Biomat-related lifestyles in the Precambrian. *PALAIOS* 14, 86–93. <http://dx.doi.org/10.2307/3515363>.
- Sen Gupta, B.K., Turner, R.E., Rabalais, N.N., 1996. Seasonal oxygen depletion in continental-shelf waters of Louisiana: historical record of benthic foraminifers. *Geology* 24, 227–230. [http://dx.doi.org/10.1130/0091-7613\(1996\)024<0227:SODICS>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1996)024<0227:SODICS>2.3.CO;2).
- Sepkoski Jr., J.J., 1979. A kinetic model of Phanerozoic taxonomic diversity II. Early Phanerozoic families and multiple equilibria. *Paleobiology* 5, 222–251. <http://dx.doi.org/10.1017/S0094837300006539>.
- Sepkoski Jr., J.J., 1981. A factor analytic description of the Phanerozoic marine fossil

- record. *Paleobiology* 7, 36–53. <http://dx.doi.org/10.1017/S0094837300003778>.
- Sepkoski Jr., J.J., 1995. The Ordovician radiations: diversification and extinction shown by global genus-level taxonomic data. In: Cooper, J.D., Droser, M.L., Finney, S.C. (Eds.), *Ordovician Odyssey*. SEPM Pacific section, Fullerton, CA, pp. 393–396 Special publication 77.
- Servais, T., Lehnert, O., Li, J.U.N., Mullins, G.L., Munnecke, A., Nützel, A., Vecoli, M., 2008. The Ordovician Biodiversification: revolution in the oceanic trophic chain. *Lethaia* 41, 99–109. <http://dx.doi.org/10.1111/j.1502-3931.2008.00115.x>.
- Servais, T., Harper, D.A.T., Munnecke, A., Owen, A.W., Sheehan, P.M., 2009. Understanding the Great Ordovician Biodiversification Event (GOBE): influences of paleogeography, paleoclimate, or paleoecology. *GSA Today* 19, 4–10. <http://dx.doi.org/10.1130/gsatg37a.1>.
- Servais, T., Owen, A.W., Harper, D.A.T., Kröger, B., Munnecke, A., 2010. The Great Ordovician Biodiversification Event (GOBE): the palaeoecological dimension. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 294, 99–119. <http://dx.doi.org/10.1016/j.palaeo.2010.05.031>.
- Servais, T., Perrier, V., Danelian, T., Klug, C., Martin, R., Munnecke, A., Nowak, H., Nützel, A., Vandenbroucke, T.R.A., Williams, M., Rasmussen, C.M.Ø., 2016. The onset of the ‘Ordovician Plankton Revolution’ in the late Cambrian. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 458, 12–28. <http://dx.doi.org/10.1016/j.palaeo.2015.11.003>.
- Shapiro, R.S., Rigby, J.K., 2004. First occurrence of an *in situ* anthaspidellid sponge in a dendrolite mound (Upper Cambrian; Great Basin, USA). *J. Paleontol.* 78, 645–650. [http://dx.doi.org/10.1666/0022-3360\(2004\)078<0645:FOOAI>2.0.CO;2](http://dx.doi.org/10.1666/0022-3360(2004)078<0645:FOOAI>2.0.CO;2).
- Sheehan, P.M., 1985. Reefs are not so different—they follow the evolutionary pattern of level-bottom communities. *Geology* 13, 46–49. [http://dx.doi.org/10.1130/0091-7613\(1985\)13<46:randsf>2.0.co;2](http://dx.doi.org/10.1130/0091-7613(1985)13<46:randsf>2.0.co;2).
- Shen, Y., Neuweiler, F., 2018. Questioning the microbial origin of automicrite in Ordovician calathid-demosponge carbonate mounds. *Sedimentology* 65, 303–333. <http://dx.doi.org/10.1111/sed.12394>.
- Shen, Y., Zhang, T., Hoffman, P.F., 2008. On the coevolution of Ediacaran oceans and animals. *Proc. Natl. Acad. Sci.* 105, 7376–7381. <http://dx.doi.org/10.1073/pnas.0802168105>.
- Shields, G.A., 2017. Earth system transition during the Tonian–Cambrian interval of biological innovation: nutrients, climate, oxygen and the marine organic carbon capacitor. In: Brasier, A.T., McLroy, D., McLoughlin, N. (Eds.), *Earth System Evolution and Early Life: A Celebration of the Work of Martin Brasier*. Geological Society of London Special Publications Vol. 448, pp. 161–177. <http://dx.doi.org/10.1144/sp448.17>.
- Shields, G.A., Carden, G.A.F., Veizer, J., Meidla, T., Rong, J.-Y., Li, R.-Y., 2003. Sr, C, and O isotope geochemistry of Ordovician brachiopods: a major isotopic event around the Middle-Late Ordovician transition. *Geochim. Cosmochim. Acta* 67, 2005–2025. [http://dx.doi.org/10.1016/S0016-7037\(02\)01116-X](http://dx.doi.org/10.1016/S0016-7037(02)01116-X).
- Signor, P.W., 1992. Taxonomic diversity and faunal turnover in the early Cambrian: did the most severe mass extinction of the Phanerozoic occur in the Botomian Stage? In: *Fifth North American Paleontological Convention Abstracts with Programs*. 6. pp. 272.
- Smith, A.B., 1990. Evolutionary diversification of echinoderms during the early Palaeozoic. In: Taylor, P.D., Larwood, G.P. (Eds.), *Major Evolutionary Radiations*. Clarendon Press, Oxford, pp. 265–286.
- Sorauf, J.E., Savarese, M., 1995. A Lower Cambrian coral from South Australia. *Palaeontology* 38, 757–770.
- Sperling, E.A., Robinson, J.M., Pisani, D., Peterson, K.J., 2010. Where's the glass? Biomarkers, molecular clocks, and microRNAs suggest a 200-Myr missing Precambrian fossil record of siliceous sponge spicules. *Geobiology* 8, 24–36. <http://dx.doi.org/10.1111/j.1472-4669.2009.00225.x>.
- Sperling, E.A., Frieder, C.A., Raman, A.V., Girguis, P.R., Levin, L.A., Knoll, A.H., 2013. Oxygen, ecology, and the Cambrian radiation of animals. *Proc. Natl. Acad. Sci.* 110, 13446–13451. <http://dx.doi.org/10.1073/pnas.1312778110>.
- Sperling, E.A., Knoll, A.H., Girguis, P.R., 2015a. The ecological physiology of Earth's second oxygen revolution. *Annu. Rev. Ecol. Syst.* 46, 215–235. <http://dx.doi.org/10.1146/annurev-ecolsys-110512-135808>.
- Sperling, E.A., Wolock, C.J., Morgan, A.S., Gill, B.C., Kunzmann, M., Halverson, G.P., Macdonald, F.A., Knoll, A.H., Johnston, D.T., 2015b. Statistical analysis of iron geochemical data suggests limited late Proterozoic oxygenation. *Nature* 523, 451–454. <http://dx.doi.org/10.1038/nature14589>.
- Spencer, B.R., 1996. The palaeoecology of some Upper Cambrian microbial-sponge-eocrinoid reefs, central Texas. In: *Sixth North American Paleontological Convention Abstracts of Papers*, pp. 367.
- Stanley, G.D., 2001. The History and Sedimentology of Ancient Reef Systems. In: *Topics in Geobiology* 17. Springer, New York (458 pp). <https://doi.org/10.1007/978-1-4615-1219-6>.
- Stearn, C.W., 2015a. Functional morphology of the Paleozoic stromatoporoid skeleton. In: Stearn, C.W. (Ed.), *Treatise on Invertebrate Paleontology, Part E, Porifera, Revised, Hypercalified Porifera, Volumes 4 and 5*. The University of Kansas Paleontological Institute, Lawrence, Kansas, pp. 551–573.
- Stearn, C.W., 2015b. Morphologic Affinities of the Paleozoic Stromatoporoidea to other fossil and recent groups. In: Stearn, C.W. (Ed.), *Treatise on Invertebrate Paleontology, Part E, Porifera, Revised, Hypercalified Porifera, Volumes 4 and 5*. The University of Kansas Paleontological Institute, Lawrence, Kansas, pp. 543–549.
- Steckbauer, A., Duarte, C.M., Carstensen, J., Vaquer-Sunyer, R., Conley, D.J., 2011. Ecosystem impacts of hypoxia: thresholds of hypoxia and pathways to recovery. *Environ. Res. Lett.* 6, 025003. <http://dx.doi.org/10.1088/1748-9326/6/2/025003>.
- Stitt, J.H., 1975. Adaptive radiation, trilobite paleoecology, and extinction, Ptychaspis Biomere, Late Cambrian of Oklahoma. In: Martinsson, A. (Ed.), *Evolution and Morphology of the Trilobita. Trilobitoidea and Merostomata, Fossils and Strata*, pp. 381–390.
- Stolley, E., 1896. Untersuchungen über coelosphaeridium, cyclocrinus, mastopora und verwandte genera des silur. In: *Archiv für Anthropologie und Geologie Schleswig-Holsteins und der benachbarten Gebiete*. 1. pp. 177–282 (in German).
- Stolper, D.A., Keller, C.B., 2018. A record of deep-ocean dissolved O₂ from the oxidation state of iron in submarine basalts. *Nature* 553, 323–327. <http://dx.doi.org/10.1038/nature25009>.
- Strauss, H., 2006. Anoxia through time. In: Neretin, L.N. (Ed.), *Past and Present Water Column Anoxia*. Springer, Amsterdam, pp. 3–19. http://dx.doi.org/10.1007/1-4020-4297-3_01.
- Sun, N., Elias, R.J., Lee, D.-J., 2014. The biological affinity of *Amsassia*: new evidence from the Ordovician of North China. *Palaeontology* 57, 1067–1089. <http://dx.doi.org/10.1111/pala.12106>.
- Taylor, J.F., 2006. History and status of the biomere concept. *Mem. Assoc. Austr. Palaeontol.* 32, 247–265.
- Taylor, M.W., Radax, R., Steger, D., Wagner, M., 2007. Sponge-associated microorganisms: evolution, ecology, and biotechnological potential. *Microbiol. Mol. Biol. Rev.* 71, 295–347. <http://dx.doi.org/10.1128/MMBR.00040-06>.
- Taylor, J.F., Repetski, J.E., Loch, J.D., Leslie, S.A., 2012. Biostratigraphy and chronostratigraphy of the Cambrian–Ordovician Great American Carbonate Bank. In: Derby, J.R., Fritz, R.D., Longacre, S.A., Morgan, W.A., Sternbach, C.A. (Eds.), *The Great American Carbonate Bank: The Geology and Economic Resources of the Cambrian–Ordovician Sauk Megasequence of Laurentia*. AAPG Memoir 98. AAPG, pp. 15–35. <http://dx.doi.org/10.1306/13331488M983497>.
- Taylor, P.D., Berning, B., Wilson, M.A., 2013. Reinterpretation of the Cambrian ‘bryozoan’ *Pywackia* as an octocoral. *J. Paleontol.* 87, 984–990. <http://dx.doi.org/10.1666/13-029>.
- Teslenko, I.L., Mambetov, A.M., Zhuravleva, I.T., Myagkova, Y.I., Meshkova, N.P., 1983. The Dedeubal bioherm belt and the history of its development. *Trudy Instituta Geologii i Geofiziki, Sibirskoye Otdeleniye* 569, 124–138 (in Russian).
- Thomas, T.R., Kavlekar, D.P., LokaBharathi, P.A., 2010. Marine drugs from sponge-microbe association—a review. *Marine Drugs* 8, 1417–1468. <http://dx.doi.org/10.3390/md8041417>.
- Thompson, C.K., Kah, L.C., 2012. Sulfur isotope evidence for widespread euxinia and a fluctuating oxycline in Early to Middle Ordovician greenhouse oceans. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 313–314, 189–214. <http://dx.doi.org/10.1016/j.palaeo.2011.10.020>.
- Thompson, C.K., Kah, L.C., Astini, R., Bowring, S.A., Buchwaldt, R., 2012. Bentonite geochronology, marine geochemistry, and the Great Ordovician Biodiversification Event (GOBE). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 321–322, 88–101. <http://dx.doi.org/10.1016/j.palaeo.2012.01.022>.
- Toomey, D.F., 1970. An unhurried look at a Lower Ordovician mound horizon, southern Franklin Mountains, west Texas. *J. Sediment. Petrol.* 40, 1318–1334. <http://dx.doi.org/10.1306/74D72194-2B21-11D7-8648000102C1865D>.
- Toomey, D.F., Ham, W.E., 1967. *Pulchrellamina*, a new mound-building organism from Lower Ordovician rocks of west Texas and southern Oklahoma. *J. Paleontol.* 41, 981–987.
- Toomey, D.F., Nitecki, M.H., 1979. Organic buildups in the Lower Ordovician (Canadian) of Texas and Oklahoma. In: *Fieldiana Geology* 2. History, Field Museum of Natural (181 pp).
- Tripathy, G.R., Hannah, J.L., Stein, H.J., Yang, G., 2014. Re-Os age and depositional environment for black shales from the Cambrian–Ordovician boundary, Green Point, western Newfoundland. *Geochim. Geophys. Geosyst.* 15, 1021–1037. <http://dx.doi.org/10.1002/gg>.
- Trotter, J.A., Williams, I.S., Barnes, C.R., Lecuyer, C., Nicoll, R.S., 2008. Did cooling oceans trigger Ordovician diversification? Evidence from conodont thermometry. *Science* 321, 550–554. <http://dx.doi.org/10.1126/science.1155814>.
- Tunnicliffe, V., 1981. High species diversity and abundance of the epibenthic community in an oxygen-deficient basin. *Nature* 294, 354–356. <http://dx.doi.org/10.1038/294354a0>.
- Tyson, R.V., Pearson, T.H., 1991. Modern and ancient continental shelf anoxia: an overview. In: Tyson, R.V., Pearson, T.H. (Eds.), *Modern and Ancient Continental Shelf Anoxia*. Geological Society of London Special Publications 58, Geological Society of London, London, pp. 1–24. <http://dx.doi.org/10.1144/GSL.SP.1991.058.01.01>.
- Van Iten, H., Marques, A.C., Leme, J.d.M., Pacheco, M.L.A.F., Simões, M.G., Smith, A., 2014. Origin and early diversification of the phylum Cnidaria Verrill: major developments in the analysis of the taxon's Proterozoic–Cambrian history. *Palaeontology* 57, 677–690. <http://dx.doi.org/10.1111/pala.12116>.
- Van Roy, P., Orr, P.J., Botting, J.P., Muir, L.A., Vinther, J., Lefebvre, B., el Hariri, K., Briggs, D.E., 2010. Ordovician faunas of Burgess Shale type. *Nature* 465, 215–218. <http://dx.doi.org/10.1038/nature09038>.
- Vinn, O., Zatoň, M., 2012. Inconsistencies in proposed annelid affinities of early biomineralized organism *Cloudina* (Ediacaran): structural and ontogenetic evidences. In: *Notebooks on Geology CG2012 A03*, pp. 39–47.
- Vologdin, A.G., 1961. Archaeocyaths and their stratigraphic significance, International Geological Congress, XX Session, Mexico. The Cambrian System, its Paleogeography and Lower Boundary Problem. Volume 3. Western Europe, Africa, USSR, Asia, America. Akademiya Nauk SSSR, Moscow, pp. 173–199.
- Vologdin, A.G., Zhuravleva, I.T., 1947. Morphology of regular archaeocyaths, Abstracts of Scientific Research for 1945. Academy of Sciences of the USSR, Division of Biological Sciences, Moscow and Leningrad, pp. 227–228.
- Wallace, M.W., Hood, A.V., Shuster, A., Greig, A., Planavsky, N.J., Reed, C.P., 2017. Oxygenation history of the Neoproterozoic to early Phanerozoic and the rise of land plants. *Earth Planet. Sci. Lett.* 466, 12–19. <http://dx.doi.org/10.1016/j.epsl.2017.02.046>.
- Webb, G.E., 1999. Youngest Early Carboniferous (late Viséan) shallow-water patch reefs in eastern Australia (Rockhampton Group, Queensland): combining quantitative micro- and macro-scale data. *Facies* 41, 111–140. <http://dx.doi.org/10.1007/BF02537462>.
- Webby, B.D., 2002. Patterns of Ordovician reef development. In: Kiessling, W., Flügel, E., Golonka, J. (Eds.), *Phanerozoic Reef Patterns*. SEPM Special Publication 72, SEPM, Tulsa, pp. 129–179. <http://dx.doi.org/10.2110/pec.02.72.0129>.
- Webby, B.D., 2015. Class uncertain, order Pulchrellaminida, new order. In: Stearn, C.W.

- (Ed.), *Treatise on Invertebrate Paleontology, Part E, Porifera, Revised, Hypercalcified Porifera*, Volumes 4 and 5. The University of Kansas Paleontological Institute, Lawrence, Kansas, pp. 837–844.
- Webby, B.D., Paris, F., Droser, M.L., Percival, I.G., 2004. The Great Ordovician Biodiversification Event. Columbia University Press, New York (497 pp).
- Werner, W., Leinfelder, R.R., Fürsich, F.T., Krautter, M., 1994. Comparative palaeoecology of marly coralline sponge-bearing reefal associations from the Kimmeridgian (Upper Jurassic) of Portugal and southwestern Germany. *Courier Forschungsinstitut Senckenberg* 172, 381–397.
- West, R.R., 2012. Evolution of the hypercalcified chaetetid-type porifera (Demospongiae). In: Stearn, C.W. (Ed.), *Treatise on Invertebrate Paleontology, Part E, Porifera, Revised, Hypercalcified Porifera*, Volumes 4 and 5. The University of Kansas Paleontological Institute, pp. 1–26.
- Westrop, S.R., Ludvigsen, R., 1987. Biogeographic control of trilobite mass extinction at an Upper Cambrian "biomere" boundary. *Paleobiology* 13, 84–99. <http://dx.doi.org/10.1017/S0094837300008605>.
- Wilde, P., Berry, W.B.N., 1984. Destabilization of the oceanic density structure and its significance to marine "extinction" events. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 48, 143–162. [http://dx.doi.org/10.1016/0031-0182\(84\)90041-5](http://dx.doi.org/10.1016/0031-0182(84)90041-5).
- Wilde, P., Quinby-Hunt, M.S., Berry, W.B.N., Orth, C.J., 1989. Palaeo-oceanography and biogeography in the Tremadoc (Ordovician) Iapetus Ocean and the origin of the chemostratigraphy of *Dictyonema flabelliforme* black shales. *Geol. Mag.* 126, 19–27. <http://dx.doi.org/10.1017/s0016756800006117>.
- Williams, M., Vannier, J., Corbari, L., Massabuau, J.C., 2011. Oxygen as a driver of early arthropod micro-benthos evolution. *PLoS One* 6, e28183. <http://dx.doi.org/10.1371/journal.pone.0028183>.
- Woo, J., 2009. The Middle Cambrian Zhangxia Formation, Shandong Province, China: depositional environments and microbial sedimentation. Seoul National University, Seoul (243 pp. Ph.D. Thesis).
- Wood, R., 1999. *Reef Evolution*. Oxford University Press, Oxford (414 pp).
- Wood, R., 2017. Palaeoecology of Ediacaran metazoan reefs. In: Brasier, A.T., McIlroy, D., McLoughlin, N. (Eds.), *Earth System Evolution and Early Life: A Celebration of the Work of Martin Brasier*. Geological Society of London Special Publications Vol. 448, pp. 195–210. <http://dx.doi.org/10.1144/sp448.1>.
- Wood, R., Erwin, D.H., 2018. Innovation not recovery: dynamic redox promotes metazoan radiations. *Biol. Rev.* 93, 863–873. <https://doi.org/10.1111/brv.12375>.
- Wood, R.A., Evans, K.R., Zhuravlev, A.Y., 1992. A new post-early Cambrian archaeocyath from Antarctica. *Geol. Mag.* 129, 491–495. <http://dx.doi.org/10.1017/S0016756800019579>.
- Wood, R., Zhuravlev, A.Y., Anaaz, C.T., 1993. The ecology of Lower Cambrian buildups from Zuune Arts, Mongolia: implications for early metazoan reef evolution. *Sedimentology* 40, 829–858. <http://dx.doi.org/10.1111/j.1365-3091.1993.tb01364.x>.
- Wood, R.A., Grotzinger, J.P., Dickson, J.A.D., 2002. Proterozoic modular biomineralized metazoan from the Nama Group, Namibia. *Science* 296, 2383–2386. <http://dx.doi.org/10.1126/science.1071599>.
- Wood, R.A., Poulton, S.W., Prave, A.R., Hoffmann, K.H., Clarkson, M.O., Guilbaud, R., Lyne, J.W., Tostevin, R., Bowyer, F., Penny, A.M., Curtis, A., Kasemann, S.A., 2015. Dynamic redox conditions control late Ediacaran metazoan ecosystems in the Nama Group, Namibia. *Precambrian Res.* 261, 252–271. <http://dx.doi.org/10.1016/j.precambres.2015.02.004>.
- Wrona, R., Zhuravlev, A.Y., 1996. Early Cambrian archaeocyaths from glacial erratics of King George Island (South Shetland Islands), Antarctica. *Palaeontol. Pol.* 55, 10–36.
- Xiao, S., 2014. Evolution: the making of Ediacaran giants. *Curr. Biol.* 24, R120–R122. <http://dx.doi.org/10.1016/j.cub.2013.12.035>.
- Yin, Z., Zhu, M., Davidson, E.H., Bottjer, D.J., Zhao, F., Tafforeau, P., 2015. Sponge grade body fossil with cellular resolution dating 60 Myr before the Cambrian. *Proc. Natl. Acad. Sci.* 112, E1453–E1460. <http://dx.doi.org/10.1073/pnas.1414577112>.
- Young, S.A., Gill, B.C., Edwards, C.T., Saltzman, M.R., Leslie, S.A., 2016. Middle–Late Ordovician (Darrivillian–Sandbian) decoupling of global sulfur and carbon cycles: isotopic evidence from eastern and southern Laurentia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 458, 118–132. <http://dx.doi.org/10.1016/j.palaeo.2015.09.040>.
- Yu, B., Dong, H., Widom, E., Chen, J., Lin, C., 2009. Geochemistry of basal Cambrian black shales and cherts from the Northern Tarim Basin, Northwest China: implications for depositional setting and tectonic history. *J. Asian Earth Sci.* 34, 418–436. <http://dx.doi.org/10.1016/j.jseas.2008.07.003>.
- Zamora, S., Clausen, S., Alvaro, J.J., Smith, A.B., 2010. Pelmatozoan echinoderms as colonizers of carbonate firmgrounds in mid-Cambrian high energy environments. *PALAIOS* 25, 764–768. <http://dx.doi.org/10.2110/palo.2010.p10-052r>.
- Zamora, S., Lefebvre, B., Javier Alvaro, J., Clausen, S., Elicki, O., Fatka, O., Jell, P., Kouchinsky, A., Lin, J.P., Nardin, E., Parsley, R., Rozhnov, S., Sprinkle, J., Sumrall, C.D., Vizcaino, D., Smith, A.B., 2013. Cambrian echinoderm diversity and palaeo-biogeography. In: Harper, D.A.T., Servais, T. (Eds.), *Early Palaeozoic Biogeography and Palaeogeography*. Geological Society, London, Memoirs 1. The Geological Society of London, London, pp. 157–171. <http://dx.doi.org/10.1144/m38.13>.
- Zamora, S., Deline, B., Javier Alvaro, J., Rahman, I.A., 2017. The Cambrian Substrate Revolution and the early evolution of attachment in suspension-feeding echinoderms. *Earth Sci. Rev.* 171, 478–491. <http://dx.doi.org/10.1016/j.earscirev.2017.06.018>.
- Zhang, X., Shu, D., Han, J., Zhang, Z., Liu, J., Fu, D., 2014. Triggers for the Cambrian explosion: hypotheses and problems. *Gondwana Res.* 25, 896–909. <http://dx.doi.org/10.1016/j.gr.2013.06.001>.
- Zhang, L., Chang, S., Khan, M.Z., Feng, Q., Danelian, T., Clausen, S., Tribouillard, N., Steiner, M., 2018. The link between metazoan diversity and paleo-oxygenation in the early Cambrian: an integrated palaeontological and geochemical record from the eastern three gorges region of South China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 495, 24–41. <http://dx.doi.org/10.1016/j.palaeo.2017.12.007>.
- Zhu, M., Babcock, L., Peng, S., 2006. Advances in Cambrian stratigraphy and paleontology: integrating correlation techniques, paleobiology, taphonomy and paleoenvironmental reconstruction. *Palaeoworld* 15, 217–222. <http://dx.doi.org/10.1016/j.palwor.2006.10.016>.
- Zhuravlev, A.Y., 1996. Reef ecosystem recovery after the Early Cambrian extinction. In: Hart, M.B. (Ed.), *Biotic Recovery from Mass Extinction Events*. Geological Society of London Special Publications 102. Geological Society of London, Oxford, pp. 79–96. <http://dx.doi.org/10.1144/GSL.SP.1996.001.01.06>.
- Zhuravlev, A.Y., 1999. A new coral from the Lower Cambrian of Siberia. *Paleontol. Zh.* 5, 27–33 (in Russian with English abstract).
- Zhuravlev, A.Y., 2001a. Biotic diversity and structure during the Neoproterozoic–Ordovician transition. In: Zhuravlev, A.Y., Riding, R. (Eds.), *Ecology of the Cambrian Radiation*. Columbia University Press, New York, pp. 173–199.
- Zhuravlev, A.Y., 2001b. Paleoeology of Cambrian reef ecosystems. In: Stanley, G.D. (Ed.), *The History and Sedimentology of Ancient Reef Systems*. Kluwer Academic/Plenum Publishers, New York, pp. 121–157. http://dx.doi.org/10.1007/978-1-4615-1219-6_4.
- Zhuravlev, A.Y., Wood, R., 1995. Lower Cambrian reefal cryptic communities. *Palaeontology* 38, 443–470.
- Zhuravlev, A.Y., Wood, R.A., 1996. Anoxia as the cause of the mid-Early Cambrian (Botomian) extinction event. *Geology* 24, 311–314. [http://dx.doi.org/10.1130/0091-7613\(1996\)024<0311:aatcot>2.3.co;2](http://dx.doi.org/10.1130/0091-7613(1996)024<0311:aatcot>2.3.co;2).
- Zhuravlev, A.Y., Debrenne, F., Lafuste, J., 1993. Early Cambrian microstructural diversification of Cnidaria. *Courier Forschungsinstitut Senckenberg* 164, 365–372.
- Zhuravlev, A.Y., Naimark, E.B., Wood, R.A., 2015. Controls on the diversity and structure of earliest metazoan communities: early Cambrian reefs from Siberia. *Earth Sci. Rev.* 147, 18–29. <http://dx.doi.org/10.1016/j.earscirev.2015.04.008>.
- Zhuravleva, I.T., Myagkova, E.I., 1987. Primitive phanerozoic multicellular organisms. In: *Institute of Geology and Geophysics, Siberian Branch, Academy of Sciences of the USSR, Transactions* 695. Nauka, Moscow (224 pp).